



**pennsylvania**  
DEPARTMENT OF HEALTH

BUREAU OF EMERGENCY MEDICAL SERVICES

# **Pennsylvania Statewide**

## **Advanced Life Support Protocols**

**Pennsylvania Department of Health  
Bureau of Emergency Medical Services**

**Effective November 1, 2008**



BUREAU OF EMERGENCY MEDICAL SERVICES

(717) 787-8740

August 1, 2008

Dear EMS Practitioner:

The Bureau of EMS, Department of Health, is pleased to provide this updated version of the Statewide ALS Protocols. The original version of the Statewide ALS Protocols was introduced on 7/1/07. New sections of the protocols that correspond to these 2008 updates are identified with **yellow highlighting**, and this will assist EMS personnel when looking for updated changes. These protocols were developed with input from hundreds of Pennsylvania's EMS stakeholders, and they were unanimously approved at a meeting of the Commonwealth's Regional EMS Medical Directors.

There are many benefits to using Statewide Protocols:

- **Uniformity** – Since many ALS practitioners provide EMS for multiple services in more than one region, statewide protocols provide a uniform expectation for EMS care. A recent report of the Institute of Medicine (IOM) has called for the development of evidence-based model pre-hospital care protocols for the treatment, triage, and transport of patients. The IOM report states that “These protocols will facilitate much more uniform treatment of injuries and illnesses across the country so that all patients will receive the current standard of care at the most appropriate location.”
- **Evidence-based** – These protocols were written to be as evidence-based as is possible. The basic pathophysiology of human illness is the same in all areas of Pennsylvania, and we would expect all patients to receive the best quality EMS care. Statewide protocols can incorporate new treatment principles, like the recent changes to the AHA guidelines, into the care in the streets on a statewide basis quickly.
- **Up-to-Date** – Regular updates to the statewide protocols will provide up-to-date protocols to all regions more rapidly than the previous process.
- **Standardization** – Standardized protocols permit better integration with the scope of practice, ALS drug list, ambulance equipment list, training curricula, and complaint investigation.

**Disaster Preparedness** – Disaster preparedness planning often crosses the boundaries of EMS regions, and the PA EMS Strike Teams have recently responded within and outside of the Commonwealth. The uniformity of statewide protocols are useful when geographically diverse EMS groups must work together to provide emergency medical care and when setting standards for disaster-related care for medical conditions that are not frequently encountered during day-to-day EMS operations.

In addition to the Statewide BLS Protocols and these Statewide ALS Protocols, regions may use additional current Department-approved regional protocols for conditions that are not covered by the Statewide BLS/ALS Protocols.

These protocols may be used by ALS personnel as soon as they are adopted by the EMS region, but all personnel must be using these protocols by the effective date of November 1, 2008. Several resources will be available to assist ALS personnel in becoming familiar with the protocol updates. These include an in-service presentation that will be available to regions and services and educational information on the Learning Management System (LMS). If you are not registered for the free LMS continuing education system, please contact the regional EMS council responsible for the area in which you live.

The Department of Health's Bureau of EMS website will always contain the most current version of the EMS protocols, the scope of practice for each level of practitioner, important EMS Information Bulletins, and many other helpful resources. This information can be accessed online at [www.health.state.pa.us/ems](http://www.health.state.pa.us/ems). The Statewide ALS Protocols may be directly printed or downloaded into a PDA for easy reference.

The Department is committed to providing Pennsylvania's EMS personnel with the most up-to-date protocols, and to do this requires periodic updates. The protocols will be reviewed annually, and EMS personnel are encouraged to provide recommendations for improvement at any time. Comments should be directed to the Commonwealth EMS Medical Director, Bureau of EMS, Rm 1034 Health & Welfare Building, 7<sup>th</sup> & Forster Streets, Harrisburg, PA 17120.

Joseph W. Schmider  
Director  
Bureau of Emergency Medical Services  
Pennsylvania Department of Health

Douglas F. Kupas, MD  
Commonwealth EMS Medical Director  
Bureau of Emergency Medical Services  
Pennsylvania Department of Health

<b>SECTION 1000:</b>	<b>Operations</b>
1000 --- General Protocol Principles .....	1000-1 thru 1000-6
<b>SECTION 2000:</b>	<b>Assessments &amp; Procedures</b>
2032 --- Confirmation of Airway Placement.....	2032-1 thru 2032-2
<b>SECTION 3000:</b>	<b>Resuscitation</b>
3001 --- Airway Obstruction.....	3001-1 thru 3001-2
3032 --- Cardiac Arrest - Traumatic.....	3032-1 thru 3032-2
3033 --- Newborn/Neonatal Resuscitation .....	3033-1 thru 3033-2
3035 --- Cardiac Arrest (Hypothermia) .....	3035-1 thru 3035-2
3041A - Ventricular Fibrillation/Pulseless VT – Adult.....	3041A-1 thru 3041A-2
3041P - Ventricular Fibrillation/Pulseless VT – Pediatric.....	3041P-1 thru 3041P-2
3042A - Asystole/Pulseless Electrical Activity (PEA) – Adult.....	3042A-1 thru 3042A-2
3042P - Asystole/Pulseless Electrical Activity (PEA) – Pediatric.....	3042P-1 thru 3042P-2
3080 --- Postresuscitation Care.....	3080-1 thru 3080-2
3091 --- Termination of Resuscitation .....	(GUIDELINE) .... 3091-1 thru 3091-2
<b>SECTION 4000:</b>	<b>Respiratory</b>
4001 --- Airway Management .....	4001-1 thru 4001-2
4002 --- Sedation-Assisted Intubation .....	(OPTIONAL) .... 4002-1 thru 4002-4
4011 --- Allergic Reaction .....	4011-1 thru 4011-2
4022 --- Asthma/COPD/Bronchospasm .....	4022-1 thru 4022-2
<b>SECTION 5000:</b>	<b>Cardiac</b>
5001 --- Suspected Acute Coronary Syndrome .....	5001-1 thru 5001-2
5002 --- Congestive Heart Failure .....	5002-1 thru 5002-2
5021A - Bradycardia - Adult .....	5021A-1 thru 5021A-2
5021P - Bradycardia - Pediatric.....	5021P-1 thru 5021P-2
5022A - Narrow Complex Tachycardia – Adult.....	5022A-1 thru 5022A-2
5022P - Narrow Complex Tachycardia – Pediatric .....	5022P-1 thru 5022P-2
5023A - Wide Complex Tachycardia – Adult.....	5023A-1 thru 5023A-2
5023P - Wide Complex Tachycardia – Pediatric.....	5023P-1 thru 5023P-2
<b>SECTION 6000:</b>	<b>Trauma &amp; Environmental</b>
6002 --- Multisystem Trauma or Traumatic Shock .....	6002-1 thru 6002-4
6003 --- Extremity Trauma.....	6003-1 thru 6003-2
6004 --- Crush Syndrome .....	6004-1 thru 6004-2
6071 --- Burns.....	6071-1 thru 6071-4
6081 --- Hypothermia / Cold Injury / Frostbite .....	6081-1 thru 6081-2
6086 --- Heat Emergencies .....	6086-1 thru 6086-2
<b>SECTION 7000:</b>	<b>Medical &amp; Ob/Gyn</b>
7002A - Altered Level of Consciousness-Adult.....	7002A-1 thru 7002A-4
7002P - Altered Level of Consciousness.....	7002P-1 thru 7002P-4
7005 --- Shock .....	7005-1 thru 7005-2
7006 --- Stroke.....	7006-1 thru 7006-4
7007 --- Seizure .....	7007-1 thru 7007-2
7009 --- Seriously Ill Appearing Patient.....	7009-1 thru 7009-2
7010 --- Nausea/Vomiting.....	7010-1 thru 7010-2
7087 --- Post-Partum Hemorrhage .....	7087-1 thru 7087-2
<b>SECTION 8000:</b>	<b>Behavioral &amp; Poisoning</b>
8001 --- Agitated Behavior/Psychiatric Disorders.....	8001-1 thru 8001-4
8031 --- Poisoning / Toxin Exposure (Ingestion / Inhalation / Absorption / Injection).....	8031-1 thru 8031-6
8081 --- Cyanide Compound Exposure .....	8081-1 thru 8081-2
8083 --- Nerve Agent/Pesticide Exposure .....	8083-1 thru 8083-4

**SECTION 9000:** **Special Considerations**

---

9001 --- Medical Command Contact..... 9001-1 thru 9001-4

---

**APPENDICES**

---

**Index..... I-1 thru I-2**

---

**GENERAL PROTOCOL PRINCIPLES  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. These general principles apply to the use of all protocols used by ALS personnel

**Purpose:**

- A. The Statewide Protocols are written with the goal of providing the highest quality of EMS patient care to patients treated by EMS practitioners in the Commonwealth.
- B. The Statewide Protocols provide a statewide uniformity and consistency to expected EMS care provided by EMS practitioners.
- C. The Statewide Protocols are written based upon the most current and best scientific evidence related to prehospital/ out-of-hospital EMS care, when this evidence is available.
- D. The Statewide Protocols are written to provide a balance between expected patient care and some educational information related to possible variations, newer information, and important warnings/ contraindications.

**Policy:****A. Scope of Practice**

1. An EMT-paramedic, with medical command authorization, may perform BLS services which may be performed by an EMT and ALS skills as defined by the EMS practitioner scope of practice as published in the PA Bulletin and listed on the EMS Bureau website when following the order of a medical command physician or when using Department approved transfer and medical treatment protocols as authorized by the ALS service medical director.
2. The Statewide BLS Protocols apply to patient care provided by ALS practitioners unless a statewide ALS protocol or Department-approved regional protocol supersedes the statewide BLS protocol.
3. The Statewide BLS Protocols and Statewide ALS Protocols apply to patient care provided by air medical services unless superseded by a Department-approved air medical service protocol.
4. Regions may establish Department-approved regional protocols, and all ALS practitioners are expected to follow applicable Department-approved regional protocols from the region in which their ALS service is based.

**B. Deviation from Protocols:**

1. When providing patient care under the EMS Act, EMS personnel must follow the orders of a medical command physician or, in the absence of such orders, the applicable protocols. In addition to the Statewide ALS Protocols, ALS practitioners must follow applicable Statewide BLS Protocols and Department-approved Regional Medical Treatment Protocols. Since written protocols cannot feasibly address all patient care situations that may develop, the Department expects EMS personnel to use their training and judgment regarding any protocol-driven care that in their judgment would be harmful to a patient under the circumstances. When the practitioner believes that following a protocol is not in the best interest of the patient, the EMS practitioner must contact a medical command physician if possible. Cases where deviation from a protocol is justified are rare. The reason for any deviation should be documented. All deviations are subject to investigation to determine whether or not they were appropriate. In all cases, EMS personnel are expected to deliver care within the scope of practice for their level of certification.
2. Medical command physicians are permitted to provide orders for patient care that are not consistent with the protocols when, under the circumstances, the procedures identified in a protocol are not the most appropriate care in the judgment of the physician or when there is not a specific protocol that is appropriate to the patient's condition. Some protocols have a section of "Possible Medical Command Orders". These are provided as a possible resource for the medical command physician and as an educational resource for the EMS personnel. These "Possible Medical Command Orders" do not substitute for the judgment of the medical command physician, and the medical command physician is under no obligation to follow the treatment options listed in this section.

3. In cases where a specific step, treatment, or medication dose within a protocol is contraindicated, EMS personnel are expected to use their judgment and training to identify these contraindications, and in these situations, the practitioner is not expected to provide that specific treatment. Failure to provide a treatment that is contraindicated is not considered a deviation from protocol, but the EMS practitioner should document the contraindication. Medical command must be contacted if the patient's condition requires alternative treatments that are not listed within the protocol.
4. Under no circumstance may a Service Medical Director institute a protocol that is separate from Department-approved Statewide or Regional protocols. Under no circumstance may a Service Medical Director institute a policy that contradicts or is not consistent with the Statewide Protocols or the Department-approved Regional Treatment Protocols.

### C. Guidelines and Protocol Options:

1. Some documents are labeled as guidelines rather than protocols. Guidelines serve as "best practice" suggestions, and these may be used by services and regions. The suggested guidelines are not considered expected care, although a region may choose to request Department approval to use a guideline as a regional protocol.
2. Some protocols or treatments within a protocol may be listed as "optional" or "if available". Regions or services may choose to use an optional protocol or treatment/medication. Regions may set requirements for options, treatments, or medications that apply to all services within the region. Service medical directors may set requirements for options, treatments, or medications that apply to all ALS practitioners within the service.  
**When statewide protocols permit options, regional requirements will supersede service level requirements regarding the options that are permitted.**
3. When protocols provide for options, regions may standardize the use of these options across the region, and services may choose options only if the choice does not conflict with regional policies.

### D. Format and Use of Protocols

1. Criteria/Exclusion Criteria - these sections list the patient conditions that are applicable to the specific protocol and list exclusion criteria that are examples of patient conditions that are not applicable to the specific protocol.
2. System Requirements - this section defines specific service or practitioner requirements that must be met in addition to the usual expectations of every EMS service or practitioner when providing treatments within the specific protocol. Most protocols are applicable to all ALS personnel, and therefore specific "system requirements" are rare.
3. Possible Medical Command Orders - this section is added for educational purposes. It provides EMS personnel with an understanding of options that may be available through medical command order, and it may be useful to medical command physicians when providing medical command orders.
4. Using the algorithm flow charts:
  - a. Although algorithms follow a step-wise approach to patient care, there are frequently several treatments that should vary in order or may be done simultaneously. Treatments that are listed within solid boxes may be done in any order, based upon the patient presentation, or may be done simultaneously when additional EMS personnel are present.
  - b. When several medication/treatment options are available, the algorithm step may refer the EMS practitioner to a "box" (outlined with a broken line) that is outside of the algorithm flow. The practitioner should refer to the box to choose the appropriate treatment and then return to the algorithm step and continue to follow the algorithm flow sheet. Regions or ALS service medical directors may define specific expectations for expected treatment options to be chosen from these boxes.
  - c. In general, the algorithms and protocols do not specify when to initiate packaging or transportation of most patients. Patient condition and paramedic judgment of the utility of on-scene treatment should determine where packaging and initiation of transport are done. If transport issues are not directly identified in the protocols, quality improvement benchmarks set by regions or service medical oversight should guide transport expectations.
  - d. EMS personnel are not required to follow every step within a protocol if a step is deemed to be inappropriate for a particular patient. For example, if a patient's condition has improved and the treatment would be unnecessary or if a medication is contraindicated.

- e. In most cases, the algorithm does not specify when or how to reassess patients. It is expected that patients are reassessed frequently, particularly after each medical intervention or medication administration. Vital signs or other appropriate reassessments should be done and documented after administering any medication that could change hemodynamic parameters, level of consciousness, etc.
  - f. Most protocols list a “Contact Medical Command” point. Although medical command should be contacted earlier if the EMS practitioner believes that consultation with a medical command physician would be helpful in treating the patient, ALS personnel must use the Medical Command Contact Protocol (# 9001), which defines actions that must be taken, when the “Contact Medical Command” point is reached.
5. Notes - these footnotes refer to the identified step of the algorithm. The notes provide additional information regarding the general step. Notes are generally used to draw attention to rarer circumstances or to provide additional educational information. Practitioners are expected to follow information within the notes as if they were a step in the algorithm flow chart.
  6. Performance Parameters - this section provides suggested benchmarks for quality improvement reviews that may occur at the service, regional or statewide level. In some instances, following quality improvement review using, at a minimum, the listed performance parameters is required.

#### **E. Use of medical command**

1. Medical command may be contacted at any step in patient care, and EMS practitioners should contact medical command if a patient’s condition is unusual and is not covered by a specific protocol, if a patient’s presentation is atypical and the protocol treatment may not be the best treatment for the patient, or in any situation where the EMS practitioner is not sure about the best treatment for the patient.
2. Service medical directors may place limitations on an EMS practitioner that require contact with medical command earlier than defined by the Statewide Protocols. These limitations may be placed upon an individual ALS practitioner when there is reason to limit the medical command authorization of that practitioner, or the limitations may apply to all service ALS practitioners for uncommon skills/procedures that may require online direction.
3. The “Medical Command Contact” Protocol # 9001 defines when medical command must be contacted and when it is appropriate to proceed beyond the “Contact Medical Command” step if communication with a medical command facility cannot be established.

#### **F. Regional and Statewide Drug Lists**

1. EMT-Paramedics may only use medications that are listed on the Statewide ALS Drug List as published in the Pennsylvania Bulletin and posted on the Bureau of EMS website.
2. Every region must publish a Department-approved regional drug list, which may not exceed the Statewide ALS Drug List. This regional list will set the standard for medications that must be carried on every ALS ambulance based within the region, and it may also list optional medications that may be carried. At a minimum, the ALS service must carry each medication that is required to provide the care that is listed in the Statewide and applicable regional protocols. This list will be used by regional council staff when conducting licensure inspections.

#### **G. Medications/Procedural Skills**

1. The protocols list many medications and treatments that are optional and are not required of every ALS Service or of every EMS practitioner. EMS regions may choose to require the use of some of these options if there is a regional reason for standardization (for example a specific medication may be required because of a regional drug box exchange program). Medications or treatments that are not required by the region may be standardized by the ALS Service Medical Director using service level policy.
2. Each region must have a Regional Drug List that includes the medications that must be carried by an ALS service and any medications that may optionally be carried by an ALS service. No medication may be listed on the Regional Drug List if it is not listed on the Statewide ALS Drug List. This list will also be used during ambulance licensure inspections.
3. General medication issues
  - a. When possible, dosing for various medications has been standardized across all protocols. EMS personnel must use their training and knowledge to assure that doses given are appropriate for the patient’s age and weight. Although doses may not exceed those listed in the protocol, it may be appropriate to decrease the doses of some medications based upon patient condition, patient vital signs or patient age.



- b. All references to medications, abbreviations, and doses have been standardized with attention to pharmacologic principles of medication error reduction.
  - c. Services should assure that medications are stored in a manner that provides for maximal shelf life and appropriate security. Some medications, for example lorazepam, may have limitations to the listed expiration date if the medication is not refrigerated. EMS services should follow Department guidance and good medication storage practices to assure that medications have not lost their potency.
  - d. EMS personnel are expected to know the contraindications for each medication and are expected to assess patients for allergies, when possible, to any medication that is given. EMS personnel should not administer medications to a patient when that medication is contraindicated in that situation.
4. Infusion mixtures - EMS regions or services may set standards for the mixture of medications that are to be given by infusion. When such standard concentrations are established, it is recommended that the region or service also provide personnel with a table to assist in administering the correct infusion dosage.
  5. Drawing blood samples - Drawing blood in the prehospital setting may assist receiving facilities in providing better diagnoses or more rapid treatment of patients, but in some areas the receiving facilities will not accept blood drawn by prehospital personnel. Although it would be appropriate for an EMS service to require blood draw in most situations where IV access is listed, EMS regions or services may determine whether drawing blood on prehospital patients is appropriate based upon the practices of local receiving hospitals.
  6. Vascular Access - Many protocols list "Initiate IV/IO NSS". The most appropriate means of establishing this peripheral vascular or intraosseous access should be determined by service policy or by the ALS practitioner's judgment based upon the condition of the patient.
    - a. Peripheral venous access may be established with a saline lock or a NSS intravenous infusion. The rate of the infusion may be KVO or should be determined by specific IV fluid volumes as stated in the appropriate protocol.
    - b. Intraosseous access - When IV access is indicated but not obtainable in a timely manner, intraosseous access is an acceptable alternative. In the absence of a regional or service policy regarding when IO access should be attempted, an IO access may be initiated based upon the ALS practitioner's judgment of the severity of the patient's condition, benefit to administration of IV fluids or medications, and proximity to the receiving facility.
      - 1) Any acceptable method or device that obtains IO access in an extremity is appropriate. Regional or service policy may indicate which technique or extremity sites are acceptable for IO access.
      - 2) Services must have the capability of providing pediatric IO access, but adult IO access is considered optional unless required by the region.
      - 3) In conscious patients, lidocaine (2%, adults= 20-40 mg, peds = 0.5 mg/kg) should be administered prior to infusing medications or fluids through an IO to reduce the pain of infusions.
    - c. Narcotic Use - when a medical command physician orders an ALS practitioner to administer a controlled substance, the medical command physician is responsible for providing a prescription in the patient's name for the order. A medical command physician may also provide a prescription for controlled substances that were given on protocol prior to contact with the medical command physician, but if the medical command physician is not comfortable writing this prescription for a medication that he/she did not order, then it is the ALS service medical director's responsibility to arrange for a prescription in the patient's name to account for the controlled substance that was administered.

#### H. Pediatric issues

1. Unless otherwise stated, pediatric protocols will apply to patients  $\leq 14$  years of age. If the patient's age is not known, then pediatric protocols will apply until there are physical signs that the patient has reached puberty/adolescence as indicated by armpit hair in boys and breast development in girls.
2. All ALS services must carry a commercial length-based device to estimate patient weight and appropriate drug dosages. When possible, these devices should be used as the primary method for determining the weight/appropriate drug doses for children. When these are not available, the following table may also be helpful:
  - a. 1 y/o 10 kg
  - b. 3 y/o 15 kg
  - c. 5 y/o 20 kg

- d. 7 y/o 25 kg
- e. 9 y/o 30 kg

### **I. Equipment Issues**

1. All medical devices must be used, maintained, and calibrated in accordance with the recommendations from the manufacturer.
2. Electronic glucose testing meters must be carried by all ALS services, and these services must have either a CLIA license or certificate of waiver. An ambulance service performing glucose testing with a meter cleared for home use by the FDA must hold a CLIA certificate of waiver. A CLIA certificate of waiver (CoW) is good for two years. Each service is responsible for determining whether a CLIA license or waiver is required.

**THIS PAGE INTENTIONALLY LEFT BLANK**

**CONFIRMATION OF AIRWAY PLACEMENT  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient who has ET tube or alternative airway device inserted by EMS personnel.

**Exclusion Criteria:**

- A. None

**System Requirements:**

- A. Every ALS ambulance service must carry and use an **electronic wave-form ETCO<sub>2</sub> detector device**<sup>1</sup> for confirmation of endotracheal tube/ alternative airway device placement.

**Procedure:**

- A. Insert ETT<sup>2</sup> or Alternative Airway Device
- B. Attach electronic ETCO<sub>2</sub> monitor to BVM.
- C. Ventilate<sup>3</sup> while simultaneously:
  - 1. Assuring “positive” CO<sub>2</sub> wave with each ventilation.
  - 2. Verifying absence of gastric sounds.
- D. Verify presence of bilateral breath sounds.
- E. Secure tube.
- F. Continuously monitor waveform ETCO<sub>2</sub>.<sup>4</sup>
- G. Reassess bilateral breath sounds and absence of gastric sounds after each move or transfer of the patient.
- H. Document all of the above.

---

**Notes:**

- 1. Colorimetric ETCO<sub>2</sub> detectors may give false negative results when the patient has had prolonged time in cardiac arrest. EDD aspiration devices may give false negative results in patients with lung disease (e.g. COPD or status asthmaticus), morbid obesity, late stages of pregnancy, or cardiac arrest. **ALS services may consider carrying colorimetric ETCO<sub>2</sub> detectors or EDD aspiration devices as back-ups in case of electronic device failure, but must primarily use the wave-form ETCO<sub>2</sub> detector as described in this procedure.**
- 2. If ETT is not visualized to pass through a good view of glottic opening, then the chance of misplaced esophageal intubation is increased and transmitted sounds during auscultation alone may lead to misdiagnosed tube position.
- 3. Immediately remove ETT or **Alternative Airway Device** if any step reveals evidence of lack of lung ventilation. If there is any doubt about adequate ventilation with an ETT or Alternative Airway Device, remove the device and ventilate with BVM.
- 4. Quantitative ETCO<sub>2</sub> readings may be beneficial in assessing the quality of CPR or as an indicator of the prognosis for successful resuscitation.


---

**Performance Parameters:**

- A. Review all ETI and **Alternative Airway Device** insertions for documentation of absence of gastric sound, presence of bilateral breath sounds, and appropriate use of a confirmation device.
- B. If systems have the capability of recording a capnograph tracing, review records of all intubated patients to assure that capnograph was recorded.
- C. Document ETCO<sub>2</sub> reading immediately after intubation, after each movement or transfer of patient and final transfer to ED stretcher.

**THIS PAGE INTENTIONALLY LEFT BLANK**

### AIRWAY OBSTRUCTION STATEWIDE ALS PROTOCOL



**AIRWAY OBSTRUCTION  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Obstructed airway from suspected foreign body.

**Exclusion Criteria:**

- A. Acute obstruction of the airway is due to systemic allergic reactions - Follow Allergic Reaction Protocol # 4011.

**Possible MC Orders:**

- A. Cricothyrotomy, if available.


---

**Notes:**

1. For children < 1 year of age, put head down and use back blows/chest thrusts. For adults and children > 1 year of age, use abdominal thrusts. For pregnant patients or patients who are too obese for abdominal thrusts, use chest thrusts.
  2. **SLAT** = **S**imultaneous **L**aryngoscopy and **A**bdominal **T**hrusts. When the foreign body can be visualized within the trachea but cannot be grasped by Magill forceps, there have been case reports of success when one rescuer visualizes the airway with a laryngoscope and another rescuer applies abdominal thrusts to temporarily dislodge the foreign body so that it can be grasped by the first rescuer with the Magill forceps
-

**CARDIAC ARREST - TRAUMATIC  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - See Protocol # 201  
Cervical spine stabilization, when indicated  
Rapid extrication



**OTHER MEDICATIONS/ TREATMENTS**  
Consider these treatments enroute if patient can arrive at a Trauma Center in < 15 minutes

**Initiate IV/IO NSS**  
**Adults:** administer NSS wide open up to 2000 mL  
**Peds:** administer NSS wide open up to 60 mL/kg

Monitor ECG / Pulse oximetry

If VF or pulseless VT, attempt defibrillation every 2 minutes (follow doses listed in VF protocols)

**Epinephrine 1:10,000;**  
**Adults:** 1 mg IV/IO 10 every 3-5 mins.  
**Peds:** 0.01 mg/kg IV/IO every 3-5 mins

**If asystole or slow PEA, administer Atropine**  
**Adults:** 1 mg IV/IO 10 every 3-5 mins  
**Peds:** 0.02 mg/kg IV/IO every 3-5 mins maximum 3 mg or 0.04 mg/kg total

Monitor ETCO<sub>2</sub>, if available

Consider other etiologies of cardiac arrest and follow appropriate protocol



**CARDIAC ARREST - TRAUMATIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient in cardiac arrest from suspected traumatic cause.

**Exclusion Criteria:**

- A. Patient that meets DOA criteria (including unwitnessed cardiac arrest of traumatic cause, decapitation, rigor mortis, etc...) – See DOA Protocol #322.
- B. Patient in cardiac arrest due to medical or non-traumatic causes <sup>1</sup>

**Possible MC Orders:**

- A. Terminate resuscitation in the field

---

**Notes:**

1. If the trauma appears to be minor and a medical condition appears to be the cause of the cardiac arrest, follow the appropriate cardiac arrest protocol.
2. If cardiac arrest is witnessed by EMS personnel, or there is evidence that the patient had any signs of life within a few minutes before the arrival of EMS personnel, proceed with this protocol. Otherwise, follow DOA Protocol # 322.
3. Unless there is an immediately correctable cause victims of traumatic cardiac arrest must arrive at a hospital within a few minutes to have any chance of survival. Placement of an advanced airway (ETT or Alternative Airway Device) or decompression of a tension pneumothorax may increase this very short time window for survival.
4. Excellent CPR is a priority:
  - a. 30 compressions: 2 ventilations in groups of 5 cycles over 2 minutes.
  - b. Push hard and fast (100/min) and allow full recoil of chest during compressions.
  - c. Change rescuer doing compressions every 2 minutes to avoid fatigue.
  - d. After advanced airway, ventilation rate should be 8-10/minute without pausing compressions to deliver ventilation.
  - e. Keep pauses in CPR to a minimum by checking rhythm when rotating rescuer doing compressions and by avoiding pauses in CPR during airway management and other interventions.
5. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032.
6. If unable to intubate on up to 3 attempts, consider alternative/ rescue airway device.
7. Transport immediately if patient can arrive at a trauma center (preferred destination) or the closest hospital in ≤ 15 minutes.
  - a. If the patient can arrive at the closest trauma center within 15 minutes, the patient should be taken to the trauma center even if another hospital is closer.
  - b. Notify the receiving facility ASAP to allow maximum time for preparation to receive the patient.
  - c. Air medical transport of patients in traumatic cardiac arrest is generally not indicated.
8. Contact medical command for possible field termination of resuscitation if the patient remains in cardiac arrest after initial resuscitation attempt and cannot arrive at the closest receiving facility within 15 minutes.
9. See Cervical Spine Immobilization Protocol # 261
10. Endotracheal medications are not very effective, but if IV/IO is unsuccessful, epinephrine, atropine, and lidocaine may be administered via endotracheal tube at twice the IV dose.
11. Field termination of resuscitation must be ordered by Medical Command Physician, otherwise continue resuscitation attempts and initiate transport.

---


**Performance Parameters:**

- A. Review all care given on scene for benefit of intervention versus potential delay to transport time. Especially procedures other than airway management and chest needle decompression in non-entrapped victims with short transport times.
- B. Review for transport to appropriate destination based upon protocol.
- C. Consider possible benchmark of on-scene time < 10 minutes for non-entrapped patients, although services may want to set goal of even less time on-scene.

**NEWBORN / NEONATAL RESUSCITATION  
STATEWIDE BLS PROTOCOL**

**BIRTH<sup>1</sup>**

Initial Patient Contact – See Protocol #201  
Consider call for second ambulance if newborn requires resuscitation



**NEWBORN / NEONATAL RESUSCITATION  
STATEWIDE BLS PROTOCOL**

**Criteria:**

- A. Newborn infant

**Exclusion Criteria:**

- A. Resuscitation may not be appropriate in rare cases where gestational age (confirmed gestational age <23 weeks) or fatal birth defects (for example anencephaly or absence of skull bones and brain hemispheres) are consistently associated with certain early death.

**Note:**

1. The newborn should be evaluated immediately after birth and reevaluated for respiratory effort, heart rate, and color every 30 seconds during the initial care until it is clear that the newborn is stable.
2. Transport the stable infant in a warm environment and within an infant car seat (if available) that has been firmly secured within the ambulance.
3. If there is evidence of meconium staining and the infant is not vigorous, use of endotracheal intubation to suction meconium from the trachea should be considered.
4. Endotracheal intubation may be appropriate at various points, although intubation is not mandatory if ventilations are positive pressure ventilations are effective. Consider ETI in the following situations:
  - a. When tracheal suctioning for meconium is required
  - b. If BVM ventilation is ineffective or prolonged
  - c. When chest compressions are performed
  - d. When endotracheal administration of medications is desired
5. Examine for central cyanosis at the face, trunk and mucous membranes. Acrocyanosis of hands and feet only is usually a normal finding if the infant is vigorous, breathing, and heart rate >100.
6. Positive pressure ventilation should use the minimum volume and pressure to achieve chest rise and /or achieve or maintain HR>100. Consider placing a gastric tube, if available, to decompress the stomach when positive pressure ventilation is required.
7. Two thumb-encircling chest technique is preferred. Compressions and ventilations should occur in a 3:1 ratio and should be done quickly enough to provide approximately 90 compressions and 30 ventilations per minute.
8. Newborns who required resuscitation are at risk for deterioration and should be transported in the environment that permits frequent reassessment.
9. After 10 minutes of continuous and adequate resuscitation efforts, discontinuation of resuscitation may be justified if there are no signs of life.

<b>APGAR SCORING CHART</b>			
<i>CLINICAL SIGNS</i>	<b>Zero</b>	<b>One</b>	<b>Two</b>
<b>A</b> = Appearance (Color)	Blue, pale	Body pink, Extremities blue	All pink
<b>P</b> = Pulse (Heart Rate)	Absent	< 100	> 100
<b>G</b> = Grimace (Reflex Response) <sup>i, ii</sup>	No response	Grimace	Cough, sneeze
<b>A</b> = Activity (Muscle Tone)	Limp	Some flexion of arms and/or legs	Well flexed
<b>R</b> = Respiratory effort	Absent	Weak cry Hypoventilation	Strong cry

<sup>i</sup>Tangential foot slap

<sup>ii</sup>Response to catheter in nostril (tested after pharynx is cleared)

**CARDIAC ARREST (HYPOTHERMIA)  
STATEWIDE ALS PROTOCOL**

**Initial Patient Contact – See Protocol #201**

- Lengthen time for breathing and pulse checks to 45 seconds each
- Begin CPR if pulseless
- Monitor ECG
- Assess body temperature, if obtainable

**If ANY breathing or pulse, Follow Hypothermia Protocol #6081**

**Follow appropriate VF/VT or Asystole/PEA protocols EXCEPT:**

- Limit defibrillation to a single attempt
- **Either:**
  - Do not administer any medications except oxygen, if temperature < 30°C (< 86°F),
  - OR*
  - Give a single dose of any medications, if temperature unknown or 30°C to 34°C (86°F to 93.2°F)
- Transport (by ground or air) ASAP to closest center capable of providing bypass rewarming <sup>1</sup>
- Notify receiving center ASAP <sup>2</sup>
- Protect against heat loss <sup>3</sup>
- Administer warmed IV/IO NSS and warmed humidified oxygen if possible

These patients may have excellent outcomes after prolonged CPR and bypass rewarming <sup>4</sup>

**TRANSPORT ASAP to Facility capable of bypass rewarming (if possible)**

**Contact Medical Command**

**If temperature > 30°C (86°F),**  
consider additional doses of medications per VF or PEA/Asystole protocol, but double the time interval between doses

**CARDIAC ARREST (HYPOTHERMIA)  
STATEWIDE ALS PROTOCOL****Criteria:**

- A.** Patient in cardiac arrest from a suspected hypothermic cause (Generalized cooling that reduces the body temperature). Hypothermia may be:
1. Acute/Immersion (e.g. sudden immersion in cold water)
  2. Subacute/Exertion (e.g. individual wandering in the woods)
  3. Chronic/ "urban" (e.g. elderly individual with no heat in home)

**Exclusion Criteria:**

- A.** Patients in cardiac arrest that meet criteria for DOA – Follow BLS DOA Protocol #322.
1. Hypothermic patient in cardiac arrest after submersion for more than 1 hour.
  2. Body tissue/chest wall frozen solid.
  3. Hypothermia patients whose body temperature has reached the temperature of the surrounding environment with other signs of death (decomposition, lividity, etc.).
- B.** Patients in cardiac arrest but without suspected hypothermia (temperature  $>34\text{ C}^{\circ}$  or  $> 92.3\text{ F}^{\circ}$ ) or who have been rewarmed to a temperature  $> 34\text{ C}^{\circ}$ , follow appropriate VF or PEA/Asystole protocol.
- C.** Patients with hypothermia (temperature  $< 34\text{ C}^{\circ}$ ) that are not in cardiac arrest. Follow Hypothermia Protocol #6081.


---

**Notes:**

1. Initiate transport to center capable of cardiac bypass rewarming (Level I and II trauma centers or other facilities known to have capability of emergency bypass rewarming) as soon as possible. Medical Command can be contacted for assistance in identifying appropriate facility and mode of transport. Consider air transport if ground transport time is  $> 30$  minutes or if it will decrease transport time. Generally air ambulances are not indicated for patients in cardiac arrest, but hypothermia is the exception to this.
  2. Notify the receiving facility as soon as possible. Bypass rewarming requires the mobilization of specialized personnel and equipment.
  3. Prevent heat loss by all means available:
    - a. Move to warm environment (like inside ambulance with heaters on maximum)
    - b. Remove wet clothing
    - c. Wrap patient in warm dry blankets
    - d. Apply heat packs to axilla, groin, and neck
  4. In severe hypothermia, EMS personnel should attempt to prevent additional heat loss, but transport should not be delayed by attempts to provide rewarming in the field.
-

**VENTRICULAR FIBRILLATION / PULSELESS VT - ADULT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - See Protocol # 201  
Assess for pulse and monitor ECG<sup>1</sup>



**VENTRICULAR FIBRILLATION / PULSELESS VT - ADULT  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Adult patient with ventricular fibrillation or pulseless ventricular tachycardia.

**Exclusion Criteria:**

- A. Cardiac arrest due to acute traumatic injury - Follow Cardiac Arrest - Traumatic Protocol #3032.
- B. Cardiac arrest due to severe hypothermia - Follow Hypothermia Protocol #3035
- C. Patient displaying an Out-of-Hospital Do Not Resuscitate (OOH-DNR) original order, bracelet, or necklace - see OOH-DNR Protocol #324.

**Possible MC Orders:**

- A. Additional antidysrhythmic therapy during cardiac arrest (magnesium sulfate 2 gm IV/ IO, procainamide 20 mg/min IV/IO if available)
- B. If tricyclic antidepressant overdose is suspected, sodium bicarbonate 1 mEq/kg IV/IO.
- C. Field termination of resuscitation.

---

**Notes:**


1. Excellent CPR is a priority:
  - a. 30 compressions: 2 ventilations in groups of 5 cycles over 2 minutes.
  - b. Push hard and fast (100/min) and allow full recoil of chest during compressions.
  - c. Change rescuer doing compressions every 2 minutes to avoid fatigue
  - d. After advanced airway, ventilation rate should be 8-10 / minute without pausing compressions to deliver ventilation. Timer on impedance threshold device (if available) or respiratory rate on ET<sub>CO</sub><sub>2</sub> monitor (if available) may help to avoid harmful hyperventilation.
  - e. Restart CPR immediately after any defibrillation attempts.
  - f. Keep pauses in CPR to a minimum by charging defibrillator during CPR, restarting compressions immediately after defibrillation without checking pulse or rhythm, and avoiding pauses in CPR during airway management.
2. Implantable Cardiac Defibrillator (ICD) may be present. Rescuer may receive slight shock, which is not dangerous.
3. If AED has been applied by BLS personnel, skip to appropriate place in protocol that incorporates previous care. ALS personnel should switch to manual defibrillator after initial AED shock.
4. Precordial thump may be used when ALS personnel witness VF arrest in a monitored patient.
5. Biphasic devices may shock at lower energy levels. Equivalent biphasic energy doses must be determined by the service AED medical director using manufacturer recommendations. Consider initial selected energy of 150-200 J for biphasic truncated exponential waveform or 120 J for rectilinear biphasic waveform. After the initial shock, subsequent shocks should be at the maximum energy available up to 360 joules.
6. Endotracheal medications are not very effective, but if IV/IO is unsuccessful, epinephrine, atropine, and lidocaine may be administered via endotracheal tube at twice the IV dose.
7. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ET<sub>CO</sub><sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032
8. If unable to intubate in up to 3 attempts, consider an alternative/ rescue airway device.
9. If available, an inspiratory impedance threshold device (ITD) may be placed on the end of an advanced airway or two-person BVM during CPR.
10. Repeat lidocaine, 0.75 mg/kg IV/IO, every 5 -10 minutes to a total dose of 3 mg/kg.
11. May repeat one additional dose of amiodarone, 150 mg IV/IO, after 10 minutes.
12. If possible, contact medical command prior to moving or transporting patient. CPR is much less effective during patient transportation, and any possible interventions by medical command will be less effective without optimal CPR.
13. Field termination of resuscitation must be ordered by Medical Command Physician, otherwise continue resuscitation attempts and initiate transport.

---

**Performance Parameters:**

- A. Documentation of ECG rhythm strips.
- B. Documentation of confirmation of advanced airway placement including documentation of gastric sounds, breath sounds and use of confirmatory device (include print out of ET<sub>CO</sub><sub>2</sub> monitor if possible)

**VENTRICULAR FIBRILLATION / PULSELESS VT - PEDIATRIC  
STATEWIDE ALS PROTOCOL**





**VENTRICULAR FIBRILLATION / PULSELESS VT - PEDIATRIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Pediatric patient (preadolescent  $\leq 14$  y/o) with ventricular fibrillation or pulseless ventricular tachycardia.

**Exclusion Criteria:**

- A. Cardiac Arrest in newborns - Follow Neonatal Resuscitation Protocol #3033.
- B. Cardiac arrest due to acute traumatic injury - Follow Cardiac Arrest - Traumatic Protocol #3032.
- C. Cardiac arrest due to severe hypothermia - Follow Cardiac Arrest - Hypothermia Protocol #3035.
- D. Patient displaying an Out-of-Hospital Do Not Resuscitate (OOH-DNR) original order, bracelet, or necklace - see OOH-DNR Protocol #324.

**Possible MC Orders:**

- A. Additional antidysrhythmic therapy during cardiac arrest (magnesium sulfate 25-50 mg/kg (max 2 gm) IV/IO, procainamide 15 mg/kg over 15-30 min IV/IO if available)
- B. If tricyclic antidepressant overdose is suspected, administer sodium bicarbonate 1-2 mEq/kg IV/IO.
- C. Field termination of resuscitation.

---

**Notes:**

1. Excellent CPR is a priority:
  - a. 15 compressions: 2 ventilations in groups of 10 cycles over 2 minutes.
  - b. Push hard and fast (100/min) and allow full recoil of chest during compressions.
  - c. Change rescuer doing compressions every 2 minutes to avoid fatigue.
  - d. After advanced airway, ventilation rate should be 8-10 / minute without pausing compressions to deliver ventilation. Respiratory rate on ETCO<sub>2</sub> monitor (if available) may help to avoid harmful hyperventilation.
  - e. Restart CPR immediately after any defibrillation attempts.
  - f. Keep pauses in CPR to a minimum by charging defibrillator during CPR, restarting compressions immediately after defibrillation without checking pulse or rhythm, and avoiding pauses in CPR during airway management.
2. If AED has been applied by BLS personnel, skip to appropriate place in protocol that incorporates previous care. ALS personnel should switch to manual defibrillator after initial AED shock.
3. Endotracheal medications are not very effective, but if IV/IO is unsuccessful, epinephrine, atropine, and lidocaine may be administered via endotracheal tube. Epinephrine 0.1 mg/kg (0.1 mL/kg of 1:1,000).
4. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032
5. Ventilation with BVM may be as effective as endotracheal intubation in children when transport times are short. If unable to intubate on up to 3 attempts, ventilate with BVM if possible.
6. Repeat lidocaine, 0.5 - 0.75 mg/kg IV, every 10 minutes to a total dose of 3 mg/kg.
7. If possible, contact medical command prior to moving or transporting patient. CPR is much less effective during patient transportation, and any possible interventions by medical command will be less effective without optimal CPR.
8. Field termination of resuscitation must be ordered by Medical Command Physician, otherwise continue resuscitation attempts and initiate transport.


---

**Performance Parameters:**

- A. Documentation of ECG rhythm strips.
- B. Documentation of confirmation of advanced airway placement including documentation of gastric sounds, breath sounds and use of confirmatory device (include print out of ETCO<sub>2</sub> monitor if possible)

**ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA) - ADULT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact- See Protocol # 201  
Monitor ECG<sup>1</sup>



**ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA) - ADULT  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Adult cardiac arrest patient presenting with asystole or organized electrical rhythm without discernable pulses.

**Exclusion Criteria:**

- A. Cardiac arrest due to acute traumatic injury- Follow Cardiac Arrest - Traumatic Protocol
- B. Cardiac arrest due to severe hypothermia - Follow Cardiac Arrest - Hypothermia Protocol #3035.
- C. Patient displaying an Out-of-Hospital Do Not Resuscitate (OOH-DNR) original order, bracelet, or necklace - Follow OOH-DNR Protocol # 324.

**Possible MC Orders:**

- A. Terminate resuscitation in the field
- B. Consider sodium bicarbonate if suspected hyperkalemia or TCA overdose.
- C. Consider calcium chloride, 10 mL of 10% solution IV (if available) if suspected renal failure/ dialysis patient or overdose of calcium channel blocker.
- D. Consider glucagon, 3-10 mg (0.05mg/kg) IV (if available) if suspected  $\beta$ -blocker overdose or calcium channel blocker overdose that is unresponsive to calcium chloride.

---

**Notes:**

1. Excellent CPR is a priority:
  - a. 30 compressions: 2 ventilations in groups of 5 cycles over 2 minutes.
  - b. Push hard and fast (100/min) and allow full recoil of chest during compressions.
  - c. Change rescuer doing compressions every 2 minutes to avoid fatigue.
  - d. After advanced airway, ventilation rate should be 8-10/minute without pausing compressions to deliver ventilation. Timer on impedance threshold device (if available) or respiratory rate on ETCO<sub>2</sub> monitor (if available) may help to avoid harmful hyperventilation.
  - e. Keep pauses in CPR to a minimum by checking rhythm when rotating rescuer doing compressions and by avoiding pauses in CPR during airway management and other interventions.
2. Confirm the presence of asystole in two leads.
3. If rhythm is unclear and possibly low-amplitude ventricular fibrillation, go to VF Protocol #3041A beginning with a full two minutes of CPR.
4. Endotracheal medications are not very effective, but if IV/IO is unsuccessful, epinephrine, atropine, and lidocaine may be administered via endotracheal tube at twice the IV dose.
5. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032.
6. If unable to intubate on up to 3 attempts, consider alternative/ rescue airway device.
7. If available, an inspiratory impedance threshold device (ITD) may be placed on the end of an advanced airway or two-person BVM during CPR.
8. If possible, contact medical command prior to moving or transporting patient. CPR is much less effective during patient transportation, and any possible interventions by medical command will be less effective without optimal CPR.
9. Field termination of resuscitation must be ordered by Medical Command Physician, otherwise continue resuscitation attempts and initiate transport.

---

**Performance Parameters:**

- A. Documentation of rhythm strips including asystole in 2 leads.
- B. Documentation of confirmation of advanced airway placement including documentation of gastric sounds, breath sounds and use of confirmatory device (include print out of ETCO<sub>2</sub> monitor if possible).

**ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA) - PEDIATRIC  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - See Protocol # 201  
Initiate CPR while monitoring for pulse  
Monitor ECG<sup>1</sup>

Asystole/ PEA

ECG<sup>2,3</sup>

VF/VT

If VF/VT, Proceed to Protocol #3041P

- Effective CPR<sup>1</sup> (15:2 for 2 rescuers) is a priority
- Every 2 minutes:
  - Rotate rescuer doing compressions and check rhythm
- Initiate IV/IO NSS
- **Epinephrine** 0.01 mg/kg IV/IO<sup>4</sup> every 3-5 mins. (0.1 mL/kg of 1:10,000)
- Manage Airway<sup>5,6</sup>
  - Ventilate 8-10 breaths/min after advanced airway
  - Consider naso/orogastric tube (if available) after patient is intubated.
- Monitor ETCO<sub>2</sub>, if available
- **Consider Other Medications/Treatments** (See Box at Right)

**OTHER MEDICATIONS/ TREATMENTS**

**If intubated, assess for tension pneumothorax or misplaced ETT:**  
If tension pneumothorax suspected, perform needle Decompression

---

**If hyperkalemia suspected in dialysis patient administer:**  
**Calcium Cl (10%)** 0.2 mL/kg IV/IO (if available)  
**Sodium bicarbonate** 1-2 mEq/kg IV/IO

---

**If hypovolemia suspected:**  
Give NSS 40 ml/kg wide open.

---

**If overdose suspected:**  
Consider **naloxone** 2 mg IV/IO

---

**If glucose < 60:**  
Administer 25% **Dextrose**, 2 mL/kg IV/IO

Change in rhythm  
(Check rhythm every 2 minutes)

YES

Proceed to Postresuscitation Protocol #3080  
OR  
Other appropriate Protocol

Contact Medical Command<sup>7</sup>

CONSIDER FIELD TERMINATION OF RESUSCITATION<sup>8</sup>  
OR  
CONTINUE RESUSCITATION AND BEGIN TRANSPORT

**ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA) - PEDIATRIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Pediatric cardiac arrest patient (preadolescent or  $\leq 14$ ) presenting with asystole or organized electrical rhythm without discernable pulses.

**Exclusion Criteria:**

- A. Cardiac arrest in newborn - Follow Neonatal Resuscitation Protocol #3033.
- B. Cardiac arrest due to acute traumatic injury - Follow Cardiac Arrest - Traumatic Protocol #3032.
- C. Cardiac arrest due to severe hypothermia - Follow Cardiac Arrest - Hypothermia Protocol #3035.
- D. Patient displaying an Out-of-Hospital Do Not Resuscitate (OOH-DNR) original order, bracelet, or necklace - Follow OOH-DNR Protocol #324.

**Possible MC Orders:**

- A. Terminate resuscitation in the field
- B. Consider sodium bicarbonate, 1-2 mEq/kg IV/IO, if suspected hyperkalemia or overdose on tricyclic antidepressant or cocaine.
- C. Consider calcium chloride, 0.2 mL/kg of 10% solution IV (if available) if suspected renal failure/dialysis patient or overdose of calcium channel blocker.
- D. Consider high-dose epinephrine, 0.1 mg/kg IV/IO (1 ml/kg of 1:10,000) if suspected  $\beta$ -blocker overdose.
- E. Consider glucagon, 0.1mg/kg IV (if available) if suspected  $\beta$ -blocker overdose or calcium channel blocker overdose that is unresponsive to calcium chloride.

---

**Notes:**

1. Excellent CPR is a priority:
  - a. 15 compressions: 2 ventilations in groups of 10 cycles over 2 minutes.
  - b. Push hard and fast (100/min) and allow full recoil of chest during compressions.
  - c. Change rescuer doing compressions every 2 minutes to avoid fatigue
  - d. After advanced airway, ventilation rate should be 8-10 / minute without pausing compressions to deliver ventilation. Respiratory rate on ETCO<sub>2</sub> monitor (if available) may help to avoid harmful hyperventilation.
  - e. Keep pauses in CPR to a minimum by avoiding pauses during airway management and other interventions.
2. Confirm the presence of asystole in two leads.
3. If rhythm is unclear and possibly low-amplitude ventricular fibrillation, then go to VF Protocol # 3041P beginning with a full two minutes of CPR.
4. Endotracheal medications are not very effective, but if IV/IO is unsuccessful, epinephrine, atropine, and lidocaine may be administered via endotracheal tube. Epinephrine 0.1 mg/kg ET ((0.1 mL/kg 1:1,000)
5. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032
6. Ventilation with BVM may be as effective as endotracheal intubation in children when transport times are short. If unable to intubate on up to 3 attempts, ventilate with BVM if possible.
7. If possible, contact medical command prior to moving or transporting patient. CPR is much less effective during patient transportation, and any possible interventions by medical command will be less effective without optimal CPR.
8. Field termination of resuscitation must be ordered by Medical Command Physician, otherwise continue resuscitation attempts and initiate transport.


---

**Performance Parameters:**

- A. Documentation of rhythm strips including asystole in 2 leads.
- B. Documentation of confirmation of advanced airway placement including documentation of gastric sounds, breath sounds and use of confirmatory device (include print out of ETCO<sub>2</sub> monitor if possible).

**POSTRESUSCITATION CARE  
STATEWIDE ALS PROTOCOL**

ROSC after cardiac arrest



**POSTRESUSCITATION CARE  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Patient that has return of spontaneous circulation (ROSC) after cardiopulmonary arrest.
  - 1. This includes resuscitation after CPR by EMS personnel and after CPR by first responders/ laypersons with or without AED use.
  - 2. The postresuscitation goals are to:
    - a. Optimize brain perfusion by optimizing cardiopulmonary function and systemic perfusion
    - b. Identify the cause/associated factors of the cardiac arrest
    - c. Prevent recurrence of cardiac arrest

**Exclusion Criteria:**

- A. Patient in cardiac arrest who does not sustain a pulse (ROSC) after resuscitation. Continue to follow appropriate cardiac arrest protocol (VF/VT, PEA/Asystole, Cardiac Arrest- Hypothermia protocols).
- B. Patients with ROSC after cardiac arrest from trauma. Continue to follow appropriate trauma protocol(s).
- C. Patient whose cardiac arrest was due to hypothermia. Follow Hypothermia protocol #6081

**Possible Medical Command Orders:**

- A. Cold (4° C) NSS bolus of 20-30 mL/kg, if available, may be ordered if patient not following commands after ROSC from nontraumatic cardiac arrest.

**Notes:**

1. If previously intubated and not tolerating endotracheal tube, administer initial dose of sedation medication. Consider extubation only if wide awake, following commands, and unable to tolerate endotracheal tube. If possible, sedation is preferred over extubation.
2. Do not permit patient to struggle against an alternative/ rescue airway. These devices should generally be removed if the patient awakens.
3. Before removing an endotracheal tube or alternative/ rescue airway device, turn patient on side and have suction running, if possible,
4. Hemodynamic instability is common after cardiac arrest, and ALS practitioners should aggressively treat hypotension to improve perfusion, especially to the brain.
5. Dobutamine drip, in addition to dopamine, may be preferable if mild hypotension (SBP 70-100) is present, while dopamine is preferred if significant hypotension or bradycardia are present. Mix dobutamine or dopamine infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP >100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**
6. Premature ventricular contractions and non-sustained VT are best treated in postresuscitation patients with oxygenation and waiting for catecholamine levels to return to normal.
7. Narrow-complex tachydysrhythmias should generally not be treated in post cardiac arrest settings unless associated with hypotension or symptoms of poor perfusion.

**POSSIBLE ASSOCIATED FACTORS:**

- Hypovolemia**  
Follow Shock protocol
- Hypoxia**  
Reassess oxygen delivery
- Hydrogen Ion (Acidosis)**  
Treat by optimizing blood pressure
- Hyperkalemia**  
Consider in dialysis patient
- Toxins**  
Follow appropriate Poisoning /Toxin protocol
- Tamponade (cardiac)**  
Follow Shock protocol
- Tension pneumothorax**  
Perform chest needle decompression, if indicated
- Thrombosis (acute MI)**  
Obtain 12-lead ECG, if available
- Trauma**  
Follow Multisystem Trauma protocol

**Sedation Options:**  
**(Choose one)**  
**(Titrate to minimum amount necessary)**

- Midazolam** 1-5 mg IV/ IO (0.05 mg/kg)  
titrated slowly  
may repeat every 5 minutes  
until maximum of 0.1 mg/kg  
*OR*
- Diazepam** 5-10 mg IV/ IO (0.1 mg/kg)  
titrated slowly  
may repeat every 5 minutes  
until maximum 0.3 mg/kg  
*OR*
- Lorazepam** 1-2 mg IV/ IO (0.1 mg/kg, max 2 mg/dose)  
titrated  
may repeat every 5 minutes  
until maximum of 4 mg

---

---

**Performance Parameters:**

- A. Review record for frequent documentation of vital signs (at least every 5 minutes for 15 minutes after cardiac arrest or for the entire time on vaspressor infusions).



**THIS PAGE LEFT INTENTIONALLY BLANK**

## TERMINATION OF RESUSCITATION STATEWIDE ALS GUIDELINE

### Purpose:

- A. When there is no response to prehospital cardiac arrest treatment, it is acceptable and often preferable to cease futile resuscitation efforts in the field.
  - 1. In patients with cardiac arrest, prehospital resuscitation is initiated with the goal of returning spontaneous circulation before permanent neurologic damage occurs. Unfortunately, most patients do not respond to an aggressive resuscitation attempt. In most situations ALS practitioners are capable of performing an initial resuscitation that is equivalent to an in-hospital resuscitation attempt, and there is usually no additional benefit to emergency department resuscitation in most cases.
  - 2. CPR that is performed during patient packaging and transport is much less effective than CPR done at the scene. Additionally, EMS personnel risk physical injury while attempting to perform CPR in a moving ambulance while unrestrained. In addition, continuing resuscitation in futile cases increases the time that EMS crews are not available for another call, impedes emergency department care of other patients, and incurs unnecessary hospital charges.
  - 3. When cardiac arrest resuscitation becomes futile, the patient's family should become the focus of the EMS personnel. Families need to be informed of what is being done, and transporting all cardiac arrest patients to the hospital is an inconvenience and inconveniences the grieving family by requiring a trip to the hospital where they must begin grieving in an unfamiliar setting. Most families understand the futility of the situation and are accepting of ceasing resuscitation efforts in the field.

### Criteria:

- A. Any cardiac arrest patient that has received resuscitation in the field but has not responded to treatment, **AND** a medical command physician has ordered termination of resuscitation efforts.
  - 1. Consider field termination of resuscitation in the following situations:
    - a. There is no response to approximately 20 minutes of ALS care including ventilation with advanced airway and several "rounds" of resuscitation drugs.
    - b. During resuscitation, new information related to DNR or terminal medical condition is obtained. If patient has OOH-DNR order, must follow OOH-DNR Protocol #324 before this protocol.
    - c. BLS care when AED has advised "no shock" on 3 sequential analyses, and the patient cannot arrive at a hospital or ALS cannot arrive at the patient within 15 minutes.

### Exclusion Criteria:

- A. Consider continuing resuscitation and transporting patients with the following conditions (although under certain circumstances, a medical command physician may order termination of resuscitation in these conditions also):
  - 1. Cardiac arrest associated with medical conditions that may have a better outcome despite prolonged resuscitation, including:
    - a. Hypothermia
    - b. Near-drowning
    - c. Lightning strike
    - d. Electrocutation
    - e. Drug overdose
  - 2. Cardiac arrest in infants and children
  - 3. Cardiac arrest in a public place
  - 4. Cardiac arrest in an environment where the bystanders do not accept the idea of ceasing efforts in the field. While most families understand the futility of the situation and are very accepting of field termination, some family members or bystanders can become hostile.

### System Requirements:

- A. Ideally, the EMS service medical director should be involved in the decision to begin a program of terminating resuscitation in the field. Each service should develop policies (e.g. related to transportation of bodies) and should make proactive contacts with key individuals (e.g. the coroner/ medical examiner, local nursing homes). Every ALS practitioner that participates in this process should have training related to "breaking bad news", dealing with grieving individuals, and interpersonal skills.

**Procedure:****A. All Patients:**

1. Follow appropriate resuscitation protocol to the point of “Contact Medical Command” to consider termination of resuscitation. Verify appropriate patient:
  - a. No femoral pulse
  - b. No respiratory efforts
  - c. Asystole or wide complex PEA at < 60 BPM
2. Contact medical command. **EMS personnel may terminate resuscitation only after order from a medical command physician.**<sup>1</sup>
3. Terminate resuscitation efforts and document time of death.
4. Consider the possibility of a crime scene. If suspected, restrict access (if possible) and notify law enforcement immediately. See Crime Scene Preservation Guideline #919.
5. Inform any family at the scene of the patient’s death and facilitate early grieving.
6. Contact the coroner or medical examiner
  - a. Do not move the body or remove any resuscitation adjuncts (e.g. endotracheal tube or IV lines) until given permission by the coroner or medical examiner.
7. Provide for dignity. If the coroner has given permission:
  - a. Remove airway devices and IV catheters
  - b. Place the patient in a position that appears comfortable
  - c. Clean up debris from the resuscitation
8. Assist the family.
  - a. Offer to call a friend, pastor, or funeral director.
  - b. Consider notifying the patient’s primary care physician.
  - c. Do not leave the scene until the family has adequate support.
9. Consider calling the local organ donation program [800-DONORS1 (Eastern PA) or 800-DONORS7 (Western PA)] for the family. Many individuals can donate corneas, skin grafts or bone grafts.
10. It is not generally the role of EMS to transport bodies, and this is usually handled by funeral directors or medical examiner offices. In some situations, EMS services may have a policy that permits transport of bodies to a local morgue for the coroner or to a local funeral director. These arrangements should not take EMS vehicles out of service for an extended time to perform these services.


---

**Notes:**

1. In remote or wilderness situations, EMS personnel **must** make every effort to contact medical command, but resuscitation may be terminated in the field without medical command when the following have occurred:
    - a. There has been no return of pulse despite >30 minutes of CPR (This does not apply in the case of hypothermia)
    - b. Transport to an emergency department will take > 30 minutes (This does not apply in the case of hypothermia)
    - c. The EMS personnel are exhausted and it is physically impossible to continue the resuscitation
-

### AIRWAY MANAGEMENT STATEWIDE ALS PROTOCOL

Assess Need for Airway or Ventilatory Support <sup>1</sup>



## AIRWAY MANAGEMENT STATEWIDE ALS PROTOCOL

**Criteria:**

- A. Any patient that requires airway management to assure adequate ventilation or a patent airway

**Exclusion Criteria:**

- A. Patient with obstructed airway- See Airway Obstruction Protocol #3001
- 

**Notes:**


1. The need for airway management is based upon the practitioner's judgment after a rapid global assessment of the patient. Indications for airway management include:
    - a. Apnea or agonal respirations
    - b. Airway reflexes compromised
    - c. Ventilatory effort compromised
    - d. Injury or medical condition compromising airway patency
    - e. Potential for future rapid compromise of airway (for example airway burns or expanding neck hematoma).
  2. If patient ventilation is initially adequate, but airway management is anticipated, high-flow oxygen should be administered. If ventilation is inadequate, provide positive pressure ventilation with high-flow oxygen (ideally, BVM ventilation should be done with two-person technique, cricoid pressure, and an oropharyngeal/ nasopharyngeal airway if possible).
  3. Techniques that may improve position for laryngoscopy are "sniffing position", head elevation, elevation of head of backboard if patient immobilized to backboard, or raising stretcher height.
  4. **Consider using external laryngeal manipulation to improve laryngoscopy view.** Consider nasotracheal intubation in patient's that are awake or have clenching of teeth. May use directional -tipped ETT or BAAM whistle to assist with nasotracheal intubation. May use bougie, lighted stylet, or fiberoptic stylet as adjuncts to endotracheal intubation (ETI).
  5. Topical atomized or nebulized lidocaine or tetracaine may be used. ALS practitioners who are qualified to perform drug-facilitated may follow the Sedation-Assisted Intubation Protocol when appropriate- See Protocol #4002. Some PHRNs may perform rapid sequence intubation when following approved air medical service protocols.
  6. Secondary/ rescue airway options may be used as the primary airway/ ventilation technique in certain situations (for example: cardiac arrest to reduce interruption in compressions, narcotic overdose until naloxone is administered, or confined/ entrapped patient in position that precludes laryngoscopy, or air medical patient inside a helicopter). Ventilation with BVM may be as effective as ETI in children when transport times are short.
  7. There should be a low threshold for using a secondary/ rescue device when basic techniques do not provide adequate ventilation, when ETI may be futile or when there have been multiple attempts at ETI.
  8. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032
  9. Placing the laryngoscope blade into the patient's mouth is considered an intubation attempt. A maximum of 3 attempts (total for all practitioners) is suggested, because the success rate dramatically decreases after 3 attempts. In some cases, it may be appropriate to proceed to a rescue airway before 3 ETI attempts have been made. Regions or service medical directors may determine the number of intubation attempts that are appropriate.
  10. ALS services must carry one type of nonsurgical Alternative/ Rescue airway device in various sizes.
- 

**Performance Parameters:**

- A. Review PCR's for documentation of the following:
1. In perfusing patients, document pulse oximetry, heart rate, and wave-form ETCO<sub>2</sub> (if available) during intubation attempts. In perfusing patients, ideally a continuous recording strip is documented.
  2. Document number of attempts at ETI and/or alternative/ rescue airway placement.
  3. Document confirmation of tube placement consistent with protocol #2032

**SEDATION-ASSISTED INTUBATION  
STATEWIDE ALS PROTOCOL [OPTIONAL]**

Assess Need for Airway or Ventilatory Support <sup>1</sup>



**SEDATION-ASSISTED INTUBATION  
STATEWIDE ALS PROTOCOL [OPTIONAL]****Sedation Options:**  
**(Choose one)****(Titrate to minimum amount necessary)****Midazolam** 1-5 mg IV/IO (0.05 mg/kg) titrated slowly  
may repeat every 5 minutes  
until maximum of 0.1 mg/kg*OR***Diazepam** 5-10 mg IV/IO (0.1 mg/kg) titrated slowly  
may repeat every 5 minutes  
until maximum 0.3 mg/kg*OR***Lorazepam** 1-2 mg IV/IO (0.1 mg/kg, max 2 mg/dose) titrated  
may repeat every 5 minutes  
until maximum of 4 mg

**SEDATION-ASSISTED INTUBATION  
STATEWIDE ALS PROTOCOL [OPTIONAL]**

**Criteria:**

- A. Sedation-assisted intubation may be appropriate for patients with compromised respiratory effort and partially intact protective airway reflexes. Examples of appropriate criteria for sedation-assisted intubation include:
1. Hypoxia (pulse oximetry < 90%) despite high flow oxygen by NRB mask or by CPAP.
  2. Inability to protect airway.
  3. Traumatic injury with GCS < 8 at the time of decision to intubate.

**Exclusion Criteria:**

- A. **CAUTION: Sedation-assisted intubation may not be appropriate for patients with fully-intact protective airway reflexes. The advantages of an airway secured by an endotracheal tube must be weighed against the potential risk of worsened hypoxia, hypotension, bradycardia, or elevated intracranial pressure that may be side effects of the sedative or complications of the intubation attempt. There is also risk of worsening a patient's outcome or misplaced esophageal intubation with this procedure. ALS practitioner judgment is critical, and practitioners must be aware of the potential adverse effects of this procedure. Other options may be preferred in some situations:**
1. CPAP and medications may be preferred if patient has acute pulmonary edema/ CHF.
  2. Nasotracheal intubation may be preferred in breathing patients.
  3. Assisting ventilation with BVM and high-flow oxygen may be preferred if ETA to receiving facility is short, if airway reflexes are fully-intact, or until naloxone can be administered in narcotic overdose.
- B. **This protocol may only be used by ALS personnel who have been approved for this skill by their service medical director and are functioning with an ALS service that meets all of the system requirements for sedation-assisted intubation. The Pennsylvania Department of Health does not condone sedation-assisted intubation by ALS practitioners or EMS services that do not meet all of the system requirements of this protocol, and does not condone the use of benzodiazepine and/or narcotic medications for the purpose of intubation when given outside of a Department approved protocol. Medical command physicians should not order such medications in an attempt to facilitate intubation.**

**System Requirements:**

- A. EMS region must approve the use of sedation-assisted intubation within the region, and the region must perform a QI audit of **every** case of sedation-assisted intubation for compliance with this protocol. All results must be forwarded to the Bureau of EMS for statewide QI.
- B. Service medical director must approve of sedation-assisted intubation by the EMS service, and must perform a QI audit of **every** case of sedation-assisted intubation for compliance with this protocol.
- C. Service medical director must personally assure training and continuing education in patient selection, endotracheal intubation, use of alternative/ rescue airway device, use of wave-form ETCO<sub>2</sub> monitoring, and use of this protocol.
- D. Service medical director must assure initial and ongoing competence (including supervised sedation-assisted intubation) for each individual EMS practitioner who will use sedation-assisted intubation. Only individuals credentialed for this procedure will perform the procedure. Medical directors should strongly consider requirements for regular supervised operating room intubations (if it is possible to arrange for such experience) and should consider the use of high-fidelity simulation as a component of assuring competence.
- E. **Two** ALS practitioners must be treating the patient before sedation-assisted intubation may be used.
- F. Service must carry an alternative/ rescue airway device in various sizes.
- G. Service must have the capability of monitoring and recording the following parameters continuously before, during and after all intubation attempts. Recordings of these parameters must be documented for every patient treated with this protocol:
1. Wave-form ETCO<sub>2</sub> (documented to confirm intubation, and monitored continuously thereafter)
  2. Heart rate by continuous ECG monitoring (documented by recording strip demonstrating trending of heart rate before, during, and after each intubation attempt).
  3. Oxygen saturation by continuous pulse oximetry (documented by recording strip demonstrating trending of pulse oximetry before, during, and after each intubation attempt).
  4. Blood pressure (documented before and immediately after intubation or intubation attempts).



- H. Etomidate may only be carried by services that follow all aspects of this protocol, and will be removed from the service's ambulances if either the service or regional QI determines that there are significant variances from this protocol.
- I. Regions or service medical directors may add more stringent criteria for use within the service. For example, services may require that medical command be contacted before sedation-assisted intubation.

---

**Notes:**

1. The need for airway management is based upon the practitioner's judgment after a rapid global assessment of the patient. Indications for airway management include:
  - a. Apnea or agonal respirations
  - b. Airway reflexes compromised
  - c. Ventilatory effort compromised
  - d. Injury or medical condition compromising airway patency
  - e. Potential for future rapid compromise of airway (for example airway burns or expanding neck hematoma).
2. If patient ventilation is initially adequate, but airway management is anticipated, high-flow oxygen should be administered. If ventilation is inadequate, provide positive pressure ventilation with high-flow oxygen (ideally, BVM ventilation should be done with two-person technique, cricoid pressure, and an oropharyngeal/ nasopharyngeal airway if possible).
3. Techniques that may improve position for laryngoscopy are "sniffing position", head elevation, elevation of head of backboard if patient immobilized to backboard, or raising stretcher height.
4. **Consider using external laryngeal manipulation to improve laryngoscopy view.** Consider nasotracheal intubation in patient's that are awake or have clenching of teeth. May use directional -tipped ETT or BAAM whistle to assist with nasotracheal intubation. May use bougie, lighted stylet, or fiberoptic stylet as adjuncts to endotracheal intubation (ETI).
5. Secondary/ rescue airway options may be used as the primary airway/ ventilation technique in certain situations (for example: cardiac arrest to reduce interruption in compressions, narcotic overdose until naloxone is administered, or confined/ entrapped patient in position that precludes laryngoscopy, or air medical patient inside a helicopter). Ventilation with BVM may be as effective as ETI in children when transport times are short.
6. There should be a low threshold for using a secondary/ rescue device when basic techniques do not provide adequate ventilation, when ETI may be futile or when there have been multiple attempts at ETI.
7. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* wave-form ETCO<sub>2</sub> monitor. Follow Confirmation of Airway Placement Protocol #2032
8. Placing the laryngoscope blade into the patient's mouth is considered an intubation attempt. A maximum of 3 attempts (total for all practitioners) is suggested, because the success rate dramatically decreases after 3 attempts. In some cases, it may be appropriate to proceed to a rescue airway before 3 ETI attempts have been made. Regions or service medical directors may determine the number of intubation attempts that are appropriate.
9. ALS services must carry one type of nonsurgical Alternative/ Rescue airway available in various sizes.

---

**Performance Parameters:**

- A. Review PCRs for documentation of the following:
  1. Review for documentation of reason for intubation.
  2. Review for complications related to intubation attempts including hypoxia, bradycardia, hypotension, and esophageal intubation(s).
  3. Review for overall successful placement of an ETT and number of attempts at ETI and alternative/ rescue airway placement.
  4. Include recording strip of continuous trend of heart rate and pulse oximetry before, during, and after each intubation attempt.
  5. Document pulse oximetry, blood pressure, and heart rate readings before and after intubation attempts. Document wave-form ETCO<sub>2</sub> readings after intubation attempts.
  6. Document number of attempts at ETI and/or alternative/ rescue airway placement.
  7. Document confirmation of tube placement by both auscultation and continuous wave-form ETCO<sub>2</sub> consistent with protocol #2032

**ALLERGIC REACTION  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - see Protocol #201  
Look for Medic Alert bracelet/necklace

Manage Airway/ Ventilate, if needed  
Apply Oxygen if needed

Monitor ECG (unless mild reaction) and Pulse Oximetry, remove stinger if visible<sup>1</sup>,  
keep part dependent if possible, apply cold pack as available

**Severe Respiratory Distress/ Wheezing or Hypotension (BP < 90 systolic)<sup>2</sup>**

**NO**

Initiate IV NSS for moderate reactions<sup>3</sup>

**Adult Patient Pediatric**

**Diphenhydramine**  
50 mg IV/IM/PO<sup>3,4</sup>

**Diphenhydramine**  
1 mg/kg IV/IM/PO<sup>3</sup>  
(max. dose 50 mg)<sup>4</sup>

**Contact Medical Command**

**YES**

**Adult Patient Pediatric**

**Epinephrine** 1:1000;  
0.3 mg IM

**Epinephrine** 1:1000;  
0.01 mg/kg IM  
(max dose 0.3 mg)

Initiate IV/IO NSS  
If Hypotension is present,  
1000 mL wide open

Initiate IV/IO NSS<sup>6</sup>  
If Hypotension is present,  
20 mL /kg wide open

**Diphenhydramine**  
50 mg IV/IO<sup>4,5</sup>

**Diphenhydramine**  
1 mg/kg IV/IO<sup>4,5</sup>  
(max. dose 50 mg)

If wheezing, Nebulized  
Bronchodilator (see box)  
May repeat continuously,  
if needed

If wheezing, Nebulized  
Bronchodilator (see box)  
May repeat  
continuously, if needed

**Contact Medical  
Command**

**Contact Medical  
Command**

Repeat **Epinephrine** IV/IO<sup>6</sup>  
Repeat IV/IO NSS bolus  
(up to 2000 mL total)  
**Methylprednisolone**  
125 mg IV/IO  
(if available)

Repeat **Epinephrine** IV/IO<sup>6</sup>  
Repeat IV/IO NSS bolus  
(up to 60 mL/kg total)  
**Methylprednisolone**  
2mg/kg IV/IO  
(if available)

**BRONCHODILATOR OPTIONS**

- **Albuterol** (approx. 2.5 mg) nebulized  
*OR*
- **Albuterol** (approx 3 mg)/ **ipratropium**  
(500 mcg) combination nebulized.  
[Half dose if ≤ 14 y/o]  
*OR*
- **Alupent** (0.2-03 mL of 0.5% diluted  
with NSS) nebulized

**ALLERGIC REACTION  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Severe Allergic Reaction/Anaphylaxis:** A patient with any of the following symptoms of severe allergic reaction after suspected exposure to an allergen (e.g. bee/wasp stings, medications/antibiotics, nuts, seafood):
1. Difficulty breathing and wheezing
  2. Difficulty breathing from swollen tongue/lips
  3. Hypotension
- B. Moderate Allergic Reaction:** A patient with less severe reaction may have:
1. Mild shortness of breath with wheezing
  2. Extensive hives and itching
  3. Mild tongue/lip swelling without difficulty swallowing or shortness of breath
- C. Mild Allergic Reaction:** A patient with a mild reaction may have:
1. Local swelling or itching isolated to extremity or area around bite site.

**Possible MC Orders:**

- A.** If unconscious or life threatening condition, consider additional doses of Epinephrine.
1. Additional dose of 1:1000 epinephrine 0.3 mg IM
  2. Epinephrine infusion (1 mg/250 mL NSS) IV/IO infused until hypotension resolves.
  3. 1:10,000 Epinephrine 0.1 mg (1mL) IV/IO very slow bolus over 5 minutes.
- B.** Glucagon, if available, (1-2 mg IV repeated every 5 minutes to 10 mg total) may be ordered if patient is taking  $\beta$ -blocker and hypotension does not resolve with NSS bolus and epinephrine.
- C.** Consider nebulized epinephrine if severe airway swelling.
- D.** Dexamethsone or hydrocortisone, if available.

**Notes:**

1. Remove stinger(s) by gently scraping stinger free with a blade or credit card, without squeezing or using forceps. In severe reaction, do not delay treatment while attempting to remove stingers.
2. In pediatrics, hypotension is  $SBP < [70 + (age \times 2)]$
3. For mild reactions, IV access is not necessary. May provide diphenhydramine, 1 mg/kg to maximum of 50 mg orally (if tablets/capsules/elixir available). May use local benzocaine applicator at bite/sting site.
4. May repeat diphenhydramine dose up to 50 mg total.
5. IV route is preferred. Diphenhydramine (Benadryl) may be given IM if IV/IO is not available.
6. Epinephrine dose may be repeated if hypotension and severe symptoms persist. Use caution when giving IV epinephrine to any patient with perfusing vital signs, especially those over 50 years old. Doses should be controlled, given slowly, and titrated only to adequate blood pressure. Higher doses may be needed in patients that are taking  $\beta$ -blocker medications. Dosing options include:
  - a. Repeat the original IM dose.
  - b. Administer dilute epinephrine infusion with frequent vital sign checks. Stop infusion when hypotension resolves:
    - i. Adults – 1 mg in 250 or 500 mL of NSS IV/IO infusion.
    - ii. Pediatrics – 0.1 mg/kg in 20 mL /kg of NSS IV/IO infusion
  - c. Administer epinephrine bolus 0.01 mg/kg (0.1 mL /kg) 1:10,000 very slowly over 10 minutes:


**Performance Parameters:**

- A.** Review for documentation of level of consciousness, airway patency, and pulse oximetry reading.

**ASTHMA / COPD / BRONCHOSPASM  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - See protocol #201

Manage Airway/ Ventilate, if needed <sup>1</sup>  
Administer Oxygen <sup>2</sup>  
Monitor Pulse Oximetry



**BRONCHODILATOR OPTIONS**

- **Albuterol** (approx. 2.5 mg) nebulized  
*OR*
- **Albuterol** (approx 3 mg)/ Ipratropium  
(500 mcg) combination nebulized.  
[Half dose if ≤ 14 y/o ]  
*OR*
- **Alupent** (0.2-03 mL of 0.5% diluted  
with NSS) nebulized

**ASTHMA / COPD / BRONCHOSPASM  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. A patient with signs and symptoms of acute respiratory distress from bronchospasm or restrictive airway disease:
  - 1. Symptoms/signs may include:
    - a. Wheezing - will have expiratory wheezing unless they are unable to move adequate air to generate wheezes
    - b. May have signs of respiratory infection (e.g. fever, nasal congestion, cough, sore throat)
    - c. May have acute onset after inhaling irritant
  - 2. This includes:
    - a. Asthma exacerbation
    - b. COPD exacerbation
    - c. Wheezing from suspected pulmonary infection (e.g. pneumonia, acute bronchitis)

**Exclusion Criteria:**

- A. Respiratory distress secondary to trauma – Follow appropriate trauma protocol.
- B. Respiratory distress secondary to congestive heart failure - Follow CHF Protocol #5002.
- C. Allergic reactions – Follow Allergic Reaction Protocol #4011.

**Possible MC Orders:**

- A. Additional nebulized bronchodilators
- B. Intravenous volume, NSS bolus or 20 mL/kg if fever, infection, or signs of dehydration.
- C. Additional doses of Epinephrine (IM or IV/IO)
- D. CPAP/BiPAP, if available and not already being used.
- E. Endotracheal Intubation, if not already done
- F. Magnesium sulfate 2 gm slow IV or infusion.

---

**Notes:**

- 1. **WARNING:** Although sometimes needed, intubation further narrows the airway restriction in a severe asthma exacerbation, and this may worsen some cases. Aggressive use of bronchodilators is generally the most important therapy for severe asthma exacerbation.
- 2. Administer oxygen at high-flow rate to all patients in severe respiratory distress. COPD patients **NOT** in respiratory distress should be given oxygen to maintain adequate O<sub>2</sub> saturation (e.g. > 90%).
- 3. Indications of severe respiratory distress include:
  - a. apprehension, anxiety, combativeness
  - b. hypoxia, SpO<sub>2</sub> < 90%
  - c. intercostals/subcostal retractions
  - d. nasal flaring
  - e. cyanosis
  - f. use of accessory muscles
- 4. **CPAP/BiPAP is required for all ALS services by 7/1/09**
- 5. Epinephrine administration may be ordered by Medical Command Physician regardless of patient's age or past medical history. Epinephrine is relatively contraindicated during pregnancy; report pregnancy to physician. Epinephrine may be repeated only with order from Medical Command Physician.


---

**Performance Parameters:**

- A. Review for documentation of lung sounds, pulse oximetry, repeat assessments/ pulse oximetry readings, and response to treatment.

### SUSPECTED ACUTE CORONARY SYNDROME STATEWIDE ALS PROTOCOL

Initial Patient Contact – see Protocol #201  
Consider non-cardiac causes <sup>1,2</sup>  
Administer Oxygen <sup>3</sup>  
Monitor Pulse Oximetry  
Monitor ECG



**SUSPECTED ACUTE CORONARY SYNDROME  
STATEWIDE ALS PROTOCOL****Criteria:**

- A.** Adult patients with symptoms of possible cardiac ischemia. Diabetics, women, and elderly patients may have atypical symptoms without retrosternal chest pain. May include:
1. Retrosternal chest heaviness/pressure/pain
  2. Radiation of pain to arm(s), neck, or jaw
  3. Associated SOB, nausea/vomiting, or sweating
  4. Possibly worsened by exertion
  5. Patient with history of recent cocaine/amphetamine use

**Exclusion Criteria:**

- A.** Chest pain/symptoms, probably not cardiac origin:
1. May include:
    - a. Pleuritic chest pain - worsens with deep breath or bending/turning
    - b. Patient less than 30 y/o

**Possible MC Orders:**

- A.** Diversion to receiving facility capable of percutaneous intervention (catheterization).
- 

**Notes:**

1. Some potentially lethal mimics of Acute Coronary Syndrome (ACS) that must be considered as the patient is assessed and treated include:
    - a. Aortic dissection
    - b. Acute pericarditis
    - c. Acute myocarditis
    - d. Spontaneous pneumothorax
    - e. Pulmonary embolism
    - f. Pneumonia/Lung infection
  2. If patient has an implanted defibrillator and is receiving shocks when not in VF/VT, may place a magnet (if available) over the AICD to deactivate it.
  3. Administer oxygen by appropriate method and monitor Pulse Oximetry. Place patient in position of comfort. Nasal cannula may be utilized if patient is unable to tolerate a facemask.
  4. Preferred method is to chew 4 baby ASA (81 mg each). Do not give aspirin if the patient has had his/her daily dose of 324 mg or has an aspirin allergy.
  5. **If a 12-lead ECG is obtained, ideally it should be transmitted to receiving/ command facility ASAP. Otherwise, give copy of all 12-lead ECGs to ED physician ASAP on arrival to facility.**
  6. If 12-lead ECG is consistent with ST-elevation MI (STEMI), either:
    - a. Follow regional destination protocol for STEMI, or
    - b. Contact medical command ASAP since some patients may benefit from transport to receiving facilities capable of percutaneous coronary interventions.
  7. Early contact with Medical Command is encouraged for patients with chest pain who have continued pain despite 3 doses of NTG, shock, or evidence of STEMI on prehospital 12-lead ECG, since these patients may benefit by direct transport to a center capable of percutaneous cardiac intervention (PCI).
  8. **WARNING:** Nitroglycerin may lead to fatal hypotension if given to patients using drugs for erectile dysfunction.
    - a. **DO NOT** administer nitroglycerin (NTG) to a patient has taken sildenafil (Viagra/Revation) or vardenafil (Levitra) within 24 hours.
    - b. **DO NOT** administer NTG to a patient who has taken tadalafil (Cialis) within the last 48 hours.
    - c. These medications may be used for conditions other than erectile dysfunction (e.g. Revation is used for pulmonary hypertension).
  9. If chest pain continues and SBP > 100, may repeat NTG approximately every 3-5 minutes as needed.
  10. If initial fluid bolus in shock protocol leads to SBP > 100, may return to this protocol and continue with NTG/analgesic medication.
- 

**Performance Parameters:**

- A.** All patients should either receive aspirin or the PCR should include documentation of why aspirin was contraindicated.
- B.** Review for appropriate transmission of 12-lead ECG when possible. Review for appropriate diversion to facility capable of PCI and/or for appropriate notification of receiving facility when STEMI is identified.
- C.** Cardiac rhythm monitored and 12-lead ECGs done (when available) and rhythm strips/12-lead ECGs documented with graphs included in PCR.
- D.** Possible benchmark for on scene time of ≤ 20 minutes.
- E.** Vital signs documented after each use of vasoactive medication (e.g. nitroglycerin or narcotic analgesics).

**CONGESTIVE HEART FAILURE  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - see Protocol #201

Manage Airway/Ventilate, if indicated


High-flow Oxygen

CPAP/BiPAP (if available – CPAP/BiPAP required for ALS by 7/1/09) if respiratory distress

AND

SpO<sub>2</sub> < 90% on High-flow Oxygen

Monitor ECG & Pulse Oximetry





**CONGESTIVE HEART FAILURE (CHF)  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patients presenting with shortness of breath from pulmonary edema/CHF, as indicated by:
1. Severe dyspnea, tachypnea, bilateral rales, tachycardia, cough with frothy sputum, or orthopnea.
  2. No fever
  3. May be associated with restlessness, agitation, pedal edema, diaphoresis, or pallor.
  4. Patient may have history of diuretic or digitalis use.

**Exclusion Criteria:**

- A. Patients presenting with shortness of breath from non-CHF etiologies:
1. Pneumonia: **WARNING** - Patients with SOB from pneumonia may have symptoms similar to those of CHF, but these patients may be harmed by diuretics. Fever may be present in these patients.
  2. COPD exacerbation: These patients may take bronchodilators without a history of diuretic use.
  3. Pneumothorax: CPAP is contraindicated in these patients.

**Possible MC Orders:**

- A. Additional Nitroglycerin
- B. Dopamine or dobutamine (if available) infusion
- C. **Captopril (if available) 25 mg sublingual or enalapril (if available) 0.625 – 1.25 mg IV**
- D. Endotracheal Intubation
- E. Morphine sulfate

---

**Notes:**

1. Relative hypotension in pulmonary edema may indicate poor cardiac function. Aggressive use of diuretics and nitroglycerin may result in extreme hypotension and further reduction of cardiac output. Contact Medical Command to discuss individualizing treatment options in these patients.
2. **WARNING:** Nitroglycerin may lead to fatal hypotension if given to patients using drugs for erectile dysfunction.
  - a. **DO NOT** give nitroglycerin (NTG) to a patient who has taken sildenafil (Viagra/Revation) or vardenafil (Levitra) within 24 hours.
  - b. **DO NOT** give NTG to a patient who has taken tadalafil (Cialis) within the last 48 hours.
  - c. These medications may be used for conditions other than erectile dysfunction (e.g. Revation is used for pulmonary hypertension).
3. **After initial single tablet/spray of NTG,** give nitroglycerin dose based upon blood pressure:
  - a. 3 SL tablets or sprays – for SBP > 180
  - b. 2 SL tablets or sprays – for SBP 140-180
  - c. 1 SL tablet or spray – for SBP 100-140
  - d. For patients on CPAP who do not tolerate SL NTG, may use 1 – 2 inches of topical nitroglycerin paste, if available.
  - e. **When available, may substitute nitroglycerin IV infusion 5 – 200 mcg / min titrated to SBP>100.**
4. NTG may be repeated every 3-5 minutes as long as blood pressure is greater than 100 systolic. **[Note: One NTG repeated every 5 minutes is equivalent to a NTG infusion of 80 mcg/min]**
5. If patient is taking prescription furosemide, administer IV dose equal to the patient's daily dose.
6. Some recommendations suggest using dobutamine for mild cardiogenic shock (SBP 70-90) and dopamine for severe shock (SBP < 70). Mix dobutamine infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP > 100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**
7. Mix dopamine infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP > 100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**

---

**Performance Parameters:**

- A. Outcomes follow-up to determine percentage of patients treated with this protocol that ultimately had hospital diagnoses of non-CHF conditions (e.g. pneumonia).
- B. Blood pressure documented after each dose of vasoactive medication (e.g. nitroglycerine)

**BRADYCARDIA – ADULT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - see Protocol # 201

**If patient has severe hypotension or impending cardiac arrest, begin Pacing IMMEDIATELY.<sup>1,2,3</sup>**

Maintain Airway/Ventilate, if needed.  
Administer Oxygen  
Monitor ECG & Pulse Oximetry  
Initiate IV/IO NSS  
Consider 12-Lead ECG, if available

**Signs or symptoms of poor perfusion?  
(e.g. acute altered mental status,  
ongoing chest pain, hypotension, or  
signs of shock)<sup>4</sup>**

NO

YES<sup>5</sup>

**Second-degree AV block  
(Type II)  
OR  
Third-degree AV block**

**Atropine 0.5 mg IV/IO<sup>6,7</sup>  
while preparing Pacer  
(May repeat if needed  
to maximum of 3 mg) OR**  
**Begin Pacing<sup>3,6,8</sup>  
Sedation<sup>3</sup>  
Initial dose, if needed  
(see box below)**

NO

YES

Consider applying  
Pacer pads<sup>5</sup>

**Pacing effective AND SBP > 100 mm/Hg**

**Observe/Monitor**

YES

NO

**CONTACT  
MEDICAL  
COMMAND**

**CONTACT  
MEDICAL  
COMMAND**

**CONTACT  
MEDICAL  
COMMAND**

Repeat additional sedation  
(see box below)

Consider **Dopamine Drip IV<sup>9</sup>**  
OR  
Consider **Epinephrine Drip<sup>10</sup>**  
Repeat additional sedation  
(see box below)

**Sedation Options<sup>3</sup>:**  
**(Choose one)**  
**(Titrate to minimum amount necessary)**  
**Midazolam** 1-5 mg IV/ IO (0.05 mg/kg) titrated slowly  
may repeat every 5 minutes  
until maximum of 0.1 mg/kg  
OR  
**Diazepam** 5-10 mg IV/ IO (0.1 mg/kg) titrated slowly  
may repeat every 5 minutes  
until maximum 0.3 mg/kg  
OR  
**Lorazepam** 1-2 mg IV/ IO (0.1 mg/kg, max 2 mg/dose) titrated  
may repeat every 5 minutes  
until maximum of 4 mg

**BRADYCARDIA - ADULT  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Adult patient with heart rate less than 60 bpm with associated symptoms.

**Exclusion Criteria:**

- A. Patient without pulse - Follow appropriate cardiac arrest protocol.
- B. History or evidence of trauma - Follow appropriate trauma protocol

**Possible MC Orders:**

- A. Additional doses of sedation or analgesia.
- B. Dopamine infusion.
- C. Glucagon 3-5 mg IV (0.05mg/kg) (if available) if beta-blocker or calcium channel blocker overdose is suspected. May be repeated in 10-15 minutes.
- D. Calcium Cl 10 mL of 10% solution IV (if available) if calcium channel-blocker overdose or hyperkalemia is suspected.

---

**Notes:**


1. When applying transcutaneous pacer for serious bradycardia or impending cardiac arrest, begin rapidly increasing the energy to obtain electrical capture.
2. Application and initiation of transcutaneous pacer should not be delayed while awaiting IV access if patient has severe symptoms.
3. Some patients may not tolerate the pacing stimulus to the skin and chest wall that occurs with transcutaneous pacing. In these cases, consider sedation if SBP > 100. (See box)
4. Consider possible etiologies:
  - a. Hyper/hypokalemia, other metabolic disorders
  - b. Hypothermia
  - c. Hypovolemia (including vomiting/diarrhea)
  - d. Hypoxia
  - e. Toxins/ overdose (e.g. beta-blocker or calcium channel-blocker)
  - f. Tamponade
  - g. Tension pneumothorax
5. Transcutaneous pacemaker electrode pads may be applied to these patients without initiating pacing so that the pacemaker is ready if patient condition deteriorates.
6. For symptomatic high-degree (second-degree or third-degree) AV block, begin pacing without delay.
7. Atropine should be administered by rapid IV push and may be repeated every 3-5 minutes, to a maximum dose of 3 mg. Atropine is ineffective and should be avoided in heart transplant patients.
8. Start pacing at heart rate of 80 and 80 mAmps. When initiating transcutaneous pacing on a patient that is conscious with a perfusing rhythm, the pacing energy level should be increased gradually to a level slightly above the minimum energy required to obtain electrical capture.
9. Mix dopamine infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP >100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**
10. Mix epinephrine infusion using regional or service prescribed concentration, and administer 2-10 mcg/min. **DO NOT exceed 10 mcg/min unless ordered by medical command physician.**

---

**Performance Parameters:**

- A. Document presence or absence of signs of poor perfusion/ shock before and after interventions.
- B. Review for appropriate use of immediate pacing before IV or atropine for patients with serious hypoperfusion or impending cardiac arrest.
- C. Documentation of correct ECG rhythm interpretation and inclusion of rhythm strips and ECGs on PCR.

### BRADYCARDIA – PEDIATRIC STATEWIDE ALS PROTOCOL



**BRADYCARDIA – PEDIATRIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Pediatric patient with heart rate < 60. Bradycardia in children is usually caused by hypoxia and often responds to oxygen and ventilatory support.

**Exclusion Criteria:**

- A. Patient without pulse - Follow appropriate cardiac arrest protocol.
- B. Newborn patient – Follow Neonatal Resuscitation Protocol #7090.
- C. History or evidence of trauma - Follow appropriate trauma protocol.
- D. Severe hypothermia – Follow Hypothermia Protocol #6081.

**Possible MC Orders:**

- A. Dopamine or epinephrine infusion.
- B. Glucagon 0.05mg/kg IV/IO (if available) if beta-blocker or calcium channel blocker overdose is suspected. May be repeated in 10-15 minutes.
- C. Calcium Cl 0.2 mL/kg of 10% solution IV/IO (if available) if calcium channel-blocker overdose or hyperkalemia is suspected.


---

**Notes:**

1. Consider possible etiologies:
    - a. Hypovolemia (including vomiting/diarrhea)
    - b. Hypoxia
    - c. Hypothermia
    - d. Hyper/hypokalemia, other metabolic disorders
    - e. Hypoglycemia
    - f. Toxins/overdose (e.g. beta-blocker or calcium channel-blocker)
    - g. Trauma/Tension Pneumothorax - follow appropriate trauma protocol.
  2. Ventilation with BVM may be as effective as endotracheal intubation in children when transport times are short. If unable to intubate on up to 3 attempts, ventilate with BVM.
  3. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032
  4. Serious signs or symptoms include:
    - a. Poor perfusion - indicated by absent or weak peripheral pulses, increased capillary refill time, skin cool/mottled.
    - b. Hypotension is SBP < 70 + (age x 2).
    - c. Respiratory difficulty (respiratory rate >60/minute) indicated by increased work of breathing (retractions, nasal flaring, grunting), cyanosis, altered level of consciousness (unusual irritability, lethargy, failure to respond to parents), stridor, wheezing.
  5. When CPR is required, a precise diagnosis of the specific bradyarrhythmia is not important. Perform chest compressions if, despite oxygenation and ventilation, the heart rate is < 60/minute and associated with poor systemic perfusion in infant or child. If severe hypothermia, do not perform chest compressions and follow Hypothermia Protocol #6081.
  6. When given IV/IO, Epinephrine may be repeated every 3-5 minutes. Epinephrine 0.1 mg/kg (1:1,000, 0.1 mL /kg) flushed with 5 mL NSS may be administered via endotracheal tube, but IV/IO route is preferred.
  7. Atropine administration may be repeated once in five minutes. Maximum dose is 1 mg per dose. Atropine 0.03 mg/kg flushed with 5 mL NSS may be administered via endotracheal tube, but IV/IO route is preferred.
-

**NARROW COMPLEX TACHYCARDIA – ADULT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact – see protocol #201



**Sedation Options:**  
(Choose one)  
(Titrate to minimum amount necessary)

**Midazolam** 1-5 mg IV/IO  
(0.05 mg/kg) titrated  
OR

**Diazepam** 5-10 mg IV/IO  
(0.1 mg/kg) titrated to effect  
OR

**Lorazepam** 1-2 mg IV/IO  
0.1 mg/kg, max 2 mg/dose) titrated

**NARROW COMPLEX TACHYCARDIA – ADULT  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Symptomatic adult patients with heart rates >100 bpm and narrow QRS complex (< 0.12 sec). It is uncommon for serious symptoms to be related to tachycardia if heart rate is <150 bpm.

**Exclusion Criteria:**

- A. Sinus tachycardia - treat underlying cause rather than rhythm. Causes may include:
  - 1. Trauma - Follow appropriate trauma protocol
  - 2. Fever
  - 3. Hypovolemia/ Shock
- B. Wide-complex tachycardias should not be treated with this protocol (SVT with wide QRS complex may be due to Wolf-Parkinson-White, and the use of calcium channel-blockers in these patients can lead to cardiac arrest.)

**Possible MC Orders:**

- A. Synchronized cardioversion
- B. Amiodarone (if available) for narrow complex irregular rhythm or for unstable patient.

**Notes:**

1. Many patients who present with SVT have evidence of cardiovascular dysfunction (low blood pressure, chest pain, congestive heart failure, altered level of consciousness). Some of these patients are unstable (such as shock, pulmonary edema, decreased level of consciousness) and require immediate synchronized cardioversion. The rest who have mild hypotension, mild shortness of breath/scattered rales, chest discomfort and a GCS > 13 may be treated with medications. If the patient develops signs/ symptoms of unstable SVT at any time during treatment, proceed immediately to the cardioversion column. The following chart illustrates the continuum from borderline to critically unstable.

<p><b><u>Borderline</u></b>                  Low BP                  SOB, Scattered Rales                  Mild chest discomfort                  Alert &amp; oriented                  GCS 14-15</p>	<p><b><u>Unstable</u></b>                  Shock                  Pulmonary Edema                  Severe chest discomfort                  Decreased level of consciousness                  GCS ≤ 13</p>
---	--

2. Regular narrow complex supraventricular tachycardias (SVTs) include reentry AV nodal tachycardia and atrial tachycardia. Atrial flutter with 2:1 conduction may be difficult to distinguish from other forms of SVT. Adenosine is not indicated if the ECG is determined to be atrial flutter or fibrillation. If atrial flutter is identified, proceed to treatment of irregular narrow complex tachycardia. If sinus tachycardia is noted, treat the underlying cause with other appropriate protocol. Fast irregular rhythms can appear regular- measure R-R intervals to be sure.
3. Avoid carotid massage if patient is older than 50 y/o or has history of hypertension.
4. Adenosine must be given by rapid IV push (over 1-3 seconds) by immediate bolus of 20 ml NSS. Adenosine success may be enhanced by administration through an antecubital IV with the arm elevated above the level of the heart during injection.
5. **Do NOT give diltiazem or verapamil if wide complex QRS or if SBP < 100.** Calcium channel blocker medications may not be the best treatment for patients with impaired ventricular function and medical command should assist with this decision.
6. May substitute verapamil 5 mg IV/IO slowly over 3-5 minutes. May repeat once at 5-10 mg dose after 15 minutes.
7. Irregular narrow complex tachycardias include atrial fibrillation, atrial flutter, or multifocal atrial tachycardia (MAT). **DO NOT** treat MAT with medications.
8. Begin with 100 joules if using a monophasic defibrillator or if ECG rhythm is atrial fibrillation.
9. If using a biphasic defibrillator, initial and subsequent countershock energy doses should be determined by service medical director.
10. Unstable patients with known chronic atrial fibrillation may be refractory to cardioversion. Consider early Medical Command contact and rapid transport.

**Performance parameters**

- A. Review for correct documentation of rhythm and for inclusion of rhythm strip in PCR
- B. Review for documentation of vital signs and rhythm after each medication or cardioversion.

**NARROW COMPLEX TACHYCARDIA – PEDIATRIC  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact – see protocol #201

Manage Airway/ Ventilate, if needed  
Apply Oxygen  
Monitor ECG & Pulse Oximetry  
Consider 12-Lead ECG, if available  
and patient stable

<p><b><u>Probable SVT</u></b></p> <ul style="list-style-type: none"> <li>• History of abrupt rate changes</li> <li>• P waves absent/abnormal</li> <li>• HR not variable</li> <li>• Infants: rate usually ≥ 220 bpm</li> <li>• Children: rate usually ≥ 180 bpm</li> </ul>	<i>OR</i>	<p><b><u>Probable Sinus Tachycardia</u></b></p> <ul style="list-style-type: none"> <li>• Known cause<sup>5</sup></li> <li>• P waves present/normal</li> <li>• Constant P-R; variable R-R</li> <li>• Infants: rate usually &lt; 220 bpm</li> <li>• Children: rate usually &lt; 180 bpm</li> </ul>
---	-----------	--

**Probable SVT**  
Unstable with signs of Poor Perfusion<sup>1</sup>

**Probable Sinus Tachycardia**

**STABLE**

**UNSTABLE<sup>1</sup>**

Assess for cause of  
sinus tachycardia<sup>5</sup>

Consider vagal  
maneuvers<sup>2</sup>  
Initiate IV/IO NSS

Initiate IV/IO NSS  
Sedation before cardioversion  
if conscious (see box at right)  
**DO NOT delay cardioversion<sup>4</sup>**  
**If IV/IO readily available<sup>4, 3</sup>**  
**Adenosine 0.1 mg/kg IV/IO<sup>3</sup>**  
**Maximum 6 mg**  
**(if available)**  
May repeat with 0.2 mg/kg IV/IO<sup>3</sup>  
**Maximum 12 mg**  
  
**If HR >180, consider**  
**Synchronized Cardioversion**  
0.5 - 1 joules/kg  
**If no conversion, repeat**  
**at 2 joules/kg,**  
**then at 4 joules/kg**

Follow other  
appropriate protocol

**Sedation Options:**  
**(Choose one)**  
**(Titrate to minimum amount necessary)**  
**Midazolam 0.05 mg/kg IV/IO**  
titrated  
*OR*  
**Diazepam 0.1 mg/kg IV/IO**  
titrated  
*OR*  
**Lorazepam 0.1 mg/kg IV/IO**  
**(max 2 mg/dose) titrated**

**Adenosine 0.1 mg/kg IV/IO<sup>3</sup>**  
Maximum 6 mg  
(if available)  
May repeat with 0.2 mg/kg IV/IO<sup>3</sup>  
Maximum 12 mg

Contact Medical Command



**NARROW COMPLEX TACHYCARDIA – PEDIATRIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Pediatric (preadolescent  $\leq$  14 years of age) patient presenting with narrow QRS complex ( $\leq$  0.08 sec) and symptomatic heart rates greater than normal for age.

**Exclusion Criteria:**

- A. Tachycardia in trauma patients (see appropriate trauma protocol)
- B. Tachycardia due to shock – Follow Shock Protocol #7005.

**Possible MC Orders:**

- A. Amiodarone (if available) 5 mg/kg IV/IO infused over 20-60 minutes.
- B. Procainamide (if available) 15 mg/kg IV/IO infused over 30-60 minutes. Avoid administering both amiodarone and procainamide.
- C. Additional synchronized cardioversions.

---

**Notes:**

1. Poor perfusion is suggested by tachypnea, altered level of consciousness, weak or absent peripheral pulses, or hypotension for age [SBP < 70 + (2 x age)].
2. Appropriate vagal maneuvers include:
  - a. Infants and young children: Cover entire face with large bag of ice without occluding the airway.
  - b. Older children: Valsalva (blow through obstructed straw) and/or carotid sinus massage.
3. Adenosine must be given by rapid IV push (over 1-3 seconds) by immediate bolus of 5 -10 mL NSS. Adenosine success may be enhanced by administration through an antecubital IV with the arm elevated above the level of the heart during injection.
4. In unstable patients, do not delay cardioversion for administration of sedation or trial of adenosine. In borderline unstable patients, adenosine may be tried and conscious patients should be sedated before cardioversion.
5. Possible causes of sinus tachycardia include:
  - a. Fever
  - b. Shock
  - c. Hypovolemia (e.g. vomiting/ diarrhea, blood loss)
  - d. Hypoxia
  - e. Abnormal electrolytes
  - f. Drug ingestions
  - g. Pneumothorax
  - h. Cardiac tamponade


---

**Performance Parameters:**

- A. Review for documentation of vital signs and rhythm after each medication or cardioversion.
- B. Review for correct documentation of rhythm and for inclusion of rhythm strip in PCR

**WIDE COMPLEX TACHYCARDIA – ADULT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact – see protocol #201



**WIDE COMPLEX TACHYCARDIA – ADULT  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Symptomatic adult patients with heart rates >100 bpm and wide QRS complex ( ≥ 0.12 sec). It is uncommon for serious symptoms to be related to tachycardia if heart rate is <150 bpm.

**Exclusion Criteria:**

- A. Sinus tachycardia with aberrancy - treat underlying cause rather than rhythm. Causes may include:
  - 1. Trauma- Follow appropriate trauma protocol
  - 2. Fever
  - 3. Hypovolemia/ Shock – Follow Shock Protocol #7005.
- B. PEA – Follow PEA Protocol #3041A.

**Possible MC Orders:**

- A. Synchronized cardioversion
- B. Amiodarone (if available) 150 mg IV/IO infused over 10 minutes. May be repeated as needed up to 2.2 gm in 24 hours.
- C. Consider sodium bicarbonate if suspected hyperkalemia or overdose.
- D. Consider calcium chloride, 10 ml of 10% solution IV (if available) if suspected renal failure/ dialysis patient or overdose of calcium channel blocker.
- E. Consider glucagon, 3-10 mg (0.05mg/kg) IV (if available) if suspected calcium channel blocker overdose that is unresponsive to calcium chloride.

---

**Notes:**

1. Many patients who present with wide complex tachycardia have evidence of cardiovascular dysfunction (low blood pressure, chest pain, congestive heart failure, altered level of consciousness). Some of these patients are unstable (such as shock, pulmonary edema, decreased level of consciousness) and require immediate synchronized cardioversion. The rest who have mild hypotension, mild shortness of breath/scattered rales, chest discomfort and a GCS >13 may be treated with medications. If the patient develops unstable signs/symptoms at any time during treatment, proceed immediately to the cardioversion column. The following chart illustrates the continuum from borderline to critically unstable.

<u>Borderline</u>	<u>Unstable</u>
Low BP	Shock
SOB, Scattered Rales	Pulmonary Edema
Mild chest discomfort	Severe chest discomfort
Alert & oriented	Decreased level of consciousness
GCS 14-15	GCS ≤ 13

2. Regular wide complex tachycardias include ventricular tachycardia and SVT with aberrancy. If the patient has a previous history of coronary artery disease, then VT is most likely. If SVT with aberrancy is suspected, adenosine (if available) may be tried. If sinus tachycardia is noted, treat the underlying cause with other appropriate protocol.
3. Vagal maneuvers may be considered. Avoid carotid massage if patient is older than 50 y/o or has history of hypertension.
4. Adenosine must be given by rapid IV push (over 1-3 seconds) by immediate bolus of 20 mL NSS. Adenosine success may be enhanced by administration through an antecubital IV with the arm elevated above the level of the heart during injection.
5. Irregular wide complex tachycardias include atrial fibrillation, pre-excitation atrial fibrillation, polymorphic VT and torsades de pointes.
6. Begin with 100 joules if using a monophasic defibrillator or if ECG rhythm is atrial fibrillation.
7. If using a biphasic defibrillator, initial and subsequent countershock energy doses should be determined by service medical director.
8. Unstable patients with known chronic atrial fibrillation may be refractory to cardioversion. Consider early Medical Command contact and rapid transport.

---

**Performance Parameters:**

- A. Review for correct documentation of rhythm and for inclusion of rhythm strip in PCR.
- B. Review for documentation of vital signs and rhythm after each medication or cardioversion.

**WIDE COMPLEX TACHYCARDIA – PEDIATRIC  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact – see protocol #201

Manage Airway/ Ventilate, if needed  
Apply Oxygen  
Monitor ECG & Pulse Oximetry  
Consider 12-Lead ECG, if available  
and patient stable

<p><b><u>Probable VT/ SVT</u></b></p> <ul style="list-style-type: none"> <li>• History of abrupt rate changes</li> <li>• P waves absent/abnormal</li> <li>• HR not variable</li> </ul>	OR	<p><b><u>Probable Sinus Tachycardia</u></b></p> <ul style="list-style-type: none"> <li>• Known cause<sup>6</sup></li> <li>• P waves present/normal</li> <li>• Constant P-R; variable R-R</li> <li>• Infants: rate usa. &lt; 220 bpm</li> <li>• Children: rate usa. &lt; 180 bpm</li> </ul>
--	----	--

**Probable VT/ SVT  
Unstable with signs of Poor Perfusion<sup>1</sup>**

**Probable Sinus Tachycardia**

**STABLE**

**UNSTABLE<sup>1</sup>**

**Assess for cause of  
sinus tachycardia<sup>6</sup>**

Consider vagal maneuvers<sup>2</sup>  
Initiate IV/IO NSS

**Contact Medical  
Command**

Initiate IV/IO NSS

**DO NOT delay cardioversion<sup>4</sup>**

**If IV/ IO readily available<sup>4, 3</sup>**  
**Adenosine 0.1 mg/kg IV/IO<sup>3</sup>**  
Maximum 6 mg (if available)  
May repeat with 0.2 mg/kg IV/IO<sup>3</sup>  
Maximum 12 mg

**Follow other  
appropriate protocol**

**Adenosine 0.1 mg/kg IV/IO<sup>3</sup>**  
Maximum 6 mg (if available)  
May repeat with 0.2 mg/kg IV/IO<sup>3</sup>  
Maximum 12 mg

**Contact Medical  
Command**

**Lidocaine 1 mg/kg IV/IO**  
OR  
**Amiodarone 5 mg/kg IV/IO**  
infused over 20-60 minutes<sup>5</sup>

**Sedation before cardioversion**  
if conscious (see box at right)

**Synchronized Cardioversion**  
**0.5 - 1 joules/kg**  
**If no conversion, repeat at**  
**2 joules/kg,**  
**then at 4 joules/kg**

**Sedation Options:  
(Choose one)  
(Titrate to minimum  
amount necessary)**

**Midazolam 0.05 mg/kg**  
IV/IO titrated  
OR  
**Diazepam 0.1 mg/kg**  
IV/IO titrated  
OR  
**Lorazepam 0.1 mg/kg**  
IV/IO (max 2 mg/dose)  
titrated

**WIDE COMPLEX TACHYCARDIA – PEDIATRIC  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Pediatric (preadolescent  $\leq$  14 years of age) patient presenting with narrow QRS complex ( $>$  0.08 sec) and symptomatic heart rates greater than normal for age

**Exclusion Criteria:**

- A. Tachycardia in trauma patients (see appropriate trauma protocol)
- B. Tachycardia due to shock – Follow Shock Protocol #7005.
- C. PEA - Follow PEA Protocol # 3041P.

**Treatment:**

- A. See accompanying flowchart.

**Possible MC Orders:**

- A. Amiodarone (if available) 5 mg/kg IV/IO infused over 20-60 minutes.
- B. Lidocaine 1 mg/kg IV/IO
- C. Procainamide (if available) 15 mg/kg IV/IO infused over 30-60 minutes. Avoid administering both amiodarone and procainamide.
- D. Additional synchronized cardioversions.
- E. Consider sodium bicarbonate, 1-2 mEq/kg IV/IO, if suspected hyperkalemia or overdose on tricyclic antidepressant or cocaine.
- F. Consider calcium chloride, 0.2 mL/kg of 10% solution IV (if available) and glucagon, 0.1 mg/kg IV/IO (if available) if suspected overdose of calcium channel blocker.
- G. **WARNING:** Calcium channel blocker medications should not be given for wide QRS rhythms.

---

**Notes:**

1. Poor perfusion is suggested by tachypnea, altered level of consciousness, weak or absent peripheral pulses, or hypotension for age [SBP  $<$  70 + (2 x age)].
2. Appropriate vagal maneuvers include:
  - a. Infants and young children: Cover entire face with large bag of ice without occluding the airway.
  - b. Older children: Valsalva (blow through obstructed straw) and/or carotid sinus massage.
3. Adenosine must be given by rapid IV push (over 1-3 seconds) by immediate bolus of 5 -10 mL NSS. Adenosine success may be enhanced by administration through an antecubital IV with the arm elevated above the level of the heart during injection.
4. In unstable patients, do not delay cardioversion for administration of sedation or trial of adenosine. In borderline unstable patients, adenosine may be tried and conscious patients should be sedated before cardioversion.
5. May substitute lidocaine, 1 mg/kg IV/IO, repeated every 5 minutes to total of 3 mg/kg.
6. Possible causes of sinus tachycardia include:
  - a. Fever
  - b. Shock
  - c. Hypovolemia (e.g. vomiting/ diarrhea, blood loss)
  - d. Hypoxia
  - e. Abnormal electrolytes
  - f. Drug ingestions
  - g. Pneumothorax
  - h. Cardiac tamponade

---

**Performance Parameters:**

- A. Review for documentation of vital signs and rhythm after each medication or cardioversion.
- B. Review for correct documentation of rhythm and for inclusion of rhythm strip in PCR.

Initial Patient Contact – See Protocol #201  
 Stabilize C-spine during assessment  
 Open airway using modified jaw thrust, if indicated.  
 Consider Air Ambulance – per Trauma Triage Protocol #180  
 Consider Rapid Extrication<sup>1</sup>

Manage Airway/Administer Oxygen/Ventilate, if needed<sup>2,3,4,5,6</sup>  
 If tension pneumothorax suspected, Needle Decompression<sup>7</sup>  
 Control External Bleeding  
 Complete Spinal Immobilization, if indicated<sup>8</sup>

**The Following Treatments Should Not Delay Transport:**

- Initiate IV/IO NSS
  - Initiate 2 large-bore IVs or single IO, if possible
  - If hypotensive, titrate NSS bolus as described<sup>9</sup>
- Monitor ECG/Pulse Oximetry
- Notify Trauma Center/receiving facility of ETA/category ASAP

**Injury Specific Treatments:**

- Follow other appropriate protocols<sup>10</sup>
- Immobilize Suspected Fractures
  - Traction splint preferred for isolated femur fracture
  - Consider pelvic binder (if available) for suspected pelvis fracture with hypotension<sup>11</sup>
- Occlude sucking chest wounds<sup>12</sup>
- Cover eviscerations<sup>13</sup>

**BEGIN TRANSPORT TO TRAUMA CENTER ASAP, if possible  
 (See Trauma Destination Protocol #180)**

**CONTACT MEDICAL COMMAND**

If hypotension persists AND due to hypovolemic shock: Repeat IV/IO NSS fluid bolus <sup>9</sup>	OR	If hypotension persists AND due to spinal cord injury: <b>Dopamine Drip<sup>14</sup></b>
--	----	---

**MULTISYSTEM TRAUMA OR TRAUMATIC SHOCK  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient that meets Category 1 or Category 2 trauma triage criteria related to traumatic injury.
- B. Patients with symptoms of spinal cord injury including extremity weakness, numbness or sensory loss.

**Exclusion Criteria:**

- A. Cardiac Arrest related to trauma – Follow BLS Cardiac Arrest – Traumatic Protocol #332.
- B. Hypotension not related to trauma – See appropriate Shock or Cardiac protocol.
- C. Patient that meets Category 3 trauma triage criteria – See appropriate injury-specific protocol.

**Possible Medical Command Orders:**

- A. Additional NSS for hypotension.
  - B. Dopamine or epinephrine infusion for neurogenic shock
  - C. Assistance with destination decisions (Trauma Center v. non-Trauma Center, Pediatric Trauma Center v. Adult Trauma Center, etc.)
- 

**Notes:**

1. Rapid extrication may be appropriate in any unsafe environment: danger of explosion (including potential secondary explosion at a terrorism incident); rapidly rising water; danger of structural collapse; hostile environments (e.g. riots); patient position prevents access to another patient that meets criteria for rapid extrication; shock; inability to establish an airway, adequately ventilate a patient, or control bleeding in entrapped position; or cardiac arrest.
2. Indications for ventilatory support include GCS < 8, inadequate respiratory effort, and airway not patent.
3. When possible, the patient should be intubated by orotracheal route using manual inline stabilization of the cervical spine. When patient's reflexes and muscle tone do not permit orotracheal intubation, consider BVM ventilation if adequate or nasotracheal intubation. Ventilation with BVM may be as effective as endotracheal intubation in children when transport times are short.
4. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds *AND* confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032.
5. If unable to intubate patient on up to 3 attempts, consider the use alternative/ rescue airway device.
6. If intubation/ventilation is needed, **AVOID OVERZEALOUS HYPERVENTILATION.**
  - a. For patients with these signs of severe head injury (GCS motor score of 1-2 or unequal/unreactive pupils), hyperventilate at:
    - 1) 20 bpm for adults
    - 2) 30 bpm for children >1 and ≤14 y/o
    - 3) 35 bpm for infant < 1 y/o
  - b. For all other trauma patients requiring ventilation, ventilate at:
    - 1) 10 bpm for adults
    - 2) 20 bpm for children > 1 and ≤ 14 y/o
    - 3) 25 bpm for infant < 1 y/o
7. Perform needle chest decompression if indicated by hypotension *AND* diminished breath sounds.
8. Follow BLS Spinal Immobilization Protocol #261.
9. IV/IO NSS fluid resuscitation should be guided by the following:
  - a. Adults: Administer NSS at wide open rate only until desired blood pressure is attained:
    - 1) When bleeding has not been controlled, titrate NSS to permit moderate hypotension (SBP between 70-90) unless severe head injury also present.
    - 2) When bleeding has been controlled or if severe head injury, titrate NSS to achieve SBP >90.
    - 3) Maximum NSS dose is 2000 mL before contacting Medical Command.
  - b. Pediatrics (preadolescent or age ≤ 14 y/o):
    - 1) When bleeding has not been controlled, titrate NSS to permit moderate hypotension (SBP between [50 + 2(age)] – [70 + 2(age)]), unless severe head injury also present.
    - 2) When bleeding has been controlled or if severe head injury, titrate NSS to achieve SBP > 70 + 2(age).
    - 3) Maximum NSS dose is 40 mL/kg before contacting Medical Command.

10. Other injury-specific appropriate protocols may include amputation, extremity trauma, burn, impaled object, or head injury.
  11. Pelvic binder splinting devices (circumferential commercial devices that compress the pelvis) are appropriate splinting devices. MAST, if available, may also be used for suspected pelvis fracture with hypotension or for suspected pelvis fracture when associated with other femur, tibia, or fibula fractures. – See MAST Suit Use Protocol #263.
  12. If sucking chest wound, cover wound with occlusive dressing sealed on 3 sides. Release dressing if worsened shortness of breath or signs of tension pneumothorax.
  13. If intestinal evisceration, cover intestines with a sterile dressing moistened with sterile saline or water; cover the area with an occlusive material (aluminum foil or plastic wrap). Cover the area with a towel or blanket to keep it warm. Transport with knees slightly flexed if possible.
    - a. **DO NOT PUSH VISCERA BACK INTO ABDOMEN**, unless prolonged extrication. In wilderness/delayed transport situations with prolonged evacuation time (at least several hours), examine the bowel for visible perforation or fecal odor. If no perforation is suspected, irrigate the eviscerated intestine with saline and gently try to replace in abdomen.
  14. Mix dopamine infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP > 100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**
- 

**Performance Parameters:**


- A. Documentation of reason for any on scene time interval over 10 minutes.
- B. Percentage of calls, without entrapment, with on scene time interval  $\leq$  10 minutes. Consider benchmark for on scene time for non-entrapped patients  $\leq$  10 minutes and  $\leq$  20 minutes for entrapped trauma patients and Category 2 trauma patients.
- C. Documentation of applicable trauma triage criteria.
- D. Appropriate destination per Trauma Patient Destination Protocol #180.
- E. Appropriate utilization of air medical transport per Trauma Patient Destination Protocol #180.



**THIS PAGE LEFT INTENTIONALLY BLANK**

**EXTREMITY TRAUMA**  
**STATEWIDE ALS PROTOCOL**

Initial Patient Contact – See Protocol #201



**Splint suspected fractures as appropriate:**

- Traction splinting is preferred over MAST for isolated femur fractures <sup>1</sup>
- Straighten severely angulated fractures if distal extremity has signs of decreased perfusion.

**If Analgesia indicated: <sup>2</sup>**

- Monitor Pulse Oximetry
- Administer oxygen, if indicated
- Initiate IV/IO NSS <sup>3</sup>
- Administer Analgesic Medication <sup>4,5</sup>  
(see box below)

**ANALGESIC MEDICATION OPTIONS**  
(Choose one)

**Fentanyl** 50-100 mcg IV/IO/**intranasal** <sup>6,7</sup> (1 mcg/kg)  
may repeat ½ dose every 5 minutes until maximum of 3 mcg/kg

*OR*

**Morphine sulfate** 2-5 mg IV <sup>6,7</sup>  
(0.05 mg/kg)  
may repeat dose every 5 minutes  
until maximum of 0.2 mg/kg

*OR*

**Nitrous Oxide** (50:50) by inhalation <sup>8</sup>

**EXTREMITY TRAUMA  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient with isolated suspected extremity fractures.
- B. Patient with extremity pain after trauma.

**Exclusion Criteria:**

- A. Multisystem trauma or traumatic/hypovolemic shock (Follow Multisystem Trauma or Traumatic Shock protocol #6002)

**Possible Medical Command Orders:**

- A. Additional fentanyl or morphine
- B. Intramuscular fentanyl or morphine

---

**Notes:**

1. Traction splinting should not be used for hip (proximal femoral neck) fractures.
  2. Narcotic pain medication should not be given if:
    - a. Oxygen saturation  $\leq$  95%
    - b. SBP  $<$  100 for adults
    - c. SBP  $<$  70 + 2(age in years) for children  $<$  14 y/o
    - d. Patient has altered level of consciousness
  3. IV/IO access is not required for administration of nitrous oxide.
  4. Narcotic pain medication may not be administered for other medical/trauma conditions (e.g. abdominal pain or multiple rib fractures) before attempted contact with Medical Command.
  5. Reassess and document 1-10 pain score after each dose of analgesia.
  6. Reduce dose for patients over 65 y/o.
  7. If respiratory depression or hypoxia occur after narcotic:
    - a. Administer oxygen and ventilate if necessary
    - b. If significant respiratory depression, administer naloxone 0.4 mg IV, titrate additional doses until adequate ventilation or total of 2 mg.
  8. Nitrous oxide should be self-administered. Patient should be coached to hold mask on his/her face, and the patient will drop mask if he/she becomes sedated. Oversedation may occur if EMS provider holds mask to patient's face. Nitrous oxide may be administered without IV access.
- 

**Performance Parameters:**

- A. Pain medication given or documentation of pain medication being offered for suspected isolated extremity fractures.
- B. Traction splinting used for isolated femur fractures without hypotension in all cases.
- C. Vital signs and oxygen saturation documented before and after any administration of narcotic.
- D. Severity of pain documented for all painful conditions, and documented before and after analgesic medications/ interventions.

**CRUSH SYNDROME  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact- See Protocol #201  
 Follow all other appropriate Trauma Protocols also  
 Administer Oxygen  
 In collapsed building, place surgical mask/ filter mask on patient, if possible <sup>1</sup>  
 Monitor ECG and pulse oximetry


Initiate IV/ IO NSS  
 Administer NSS bolus of 20 mL /kg (warm if possible) <sup>2,3</sup>

**Patient Entrapped?**

**NO**

See Multisystem Trauma Protocol #6002, Extremity Injury Protocol #6003, or other appropriate protocol(s)

**YES**



**CRUSH SYNDROME  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient entrapped and crushed under heavy load (e.g. extremities and/or body crushed in building collapse, trench collapse, industrial accident, or pinned under/by heavy vehicle/ farm equipment for more than 30 minutes and with suspected crush syndrome:

**Exclusion Criteria:**

- A. Patient trapped for less than 30 minutes
- B. Patient entrapped but without significant tissue crushing.
- C. Altered mental status – See Altered Level of Consciousness Protocol #7002A or #7002P.

**Possible Medical Command Orders:**

- A. Additional narcotic analgesic
- B. Alteration in intravenous fluid volume
- C. Benzodiazepine for anxiety
- D. Additional sodium bicarbonate

**Notes:**

1. In addition to a dust mask, victims may need eye, head and hearing protection if rescue crews are working nearby.
2. Avoid using Ringer's lactate as an IV fluid due to its potassium content.
3. Victims of structural collapse may become hypothermic in prolonged incidents, even in warm weather.
4. Pain control is important; crush injury is painful, often increasing upon release.
5. EKG signs of severe and life threatening hyperkalemia should be treated with calcium chloride and sodium bicarbonate using dose in protocol.
  - a. Signs of hyperkalemia:
    - 1) Mild – peaked T waves
    - 2) Moderate – prolonged PR interval, decreased P wave amplitude, ST segment depression
    - 3) Severe – QRS widening, flat P waves, second degree Mobitz I, PVCs
    - 4) Life Threatening – absent P waves, V-tach, AV block, extremely widened QRS

**Performance Parameters:**

- A. Review every case where crush syndrome protocol used.

**BURNS  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - Protocol #201  
Use PPE/Remove from source of burn <sup>1</sup>  
Follow BLS Burn Protocol # 671

**History/Evidence of  
Category 1 or 2 Trauma**

YES →

**Proceed to  
Multisystem  
Trauma Protocol  
#6001**

NO ↓

Manage Airway/Ventilate, if needed <sup>2,3,4</sup>  
Administer Oxygen, if indicated <sup>2</sup>  
Immobilize Spine, as indicated by BLS Protocol #261

<b>Mechanism of burn injury</b>		
<b><u>Chemical</u></b> Brush off dry, then flush with water <sup>5</sup>	<b><u>Thermal</u></b> Dry, sterile sheet Cool, unless large BSA involved	<b><u>Electrical / Lightning</u></b> Monitor ECG Dry, sterile dressing to entrance and exit wounds

Monitor Pulse Oximetry and/or ECG, if indicated <sup>2,6</sup>  
Determine Burn Extent & Severity <sup>7</sup> (rule of nines)  
Initiate IV/IO NSS, if indicated

Administer 20 mL /kg NSS wide open for  
hypotension or severe burn, <sup>8</sup>  
Administer Analgesic Medication (see box below),  
if indicated <sup>9</sup>

**TRANSPORT TO CLOSEST  
APPROPRIATE FACILITY/  
TRAUMA CENTER <sup>13</sup>**

**Contact Medical Command <sup>14</sup>**

If hypotension persists or if extensive BSA burn,  
repeat 20 mL /kg NSS fluid bolus <sup>8</sup>  
If pain continues, Administer Repeat dose(s) of Analgesic Medication  
(see box below)

**ANALGESIC MEDICATION OPTIONS**  
(Choose one)

**Fentanyl** 50-100 mcg IV/IO/**intranasal** <sup>10,11</sup> (1 mcg/kg)  
may repeat ½ dose every 5 minutes until maximum of 3 mcg/kg  
*OR*  
**Morphine sulfate** 2-5 mg IV <sup>10,11</sup> (0.05 mg/kg)  
may repeat dose every 5 minutes until maximum of 0.2 mg/kg  
*OR*  
**Nitrous Oxide** (50:50) by inhalation <sup>12</sup>

**BURNS**  
**STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Patient with burns from:
  - 1. Thermal injury
  - 2. Chemical dermal injury.
- B. Patient with lightning or electrical injury.

**Possible MC Orders:**

- A. Additional morphine or fentanyl
- B. Transport to a burn center or trauma center
- C. CPAP/BiPAP for respiratory difficulty

---

**Notes:**

1. Consider scene safety. Be aware of possible chemical contamination and/or electrical sources. Stop the burning process. Remove clothing and constricting jewelry.
2. Determine presence of respiratory burns as indicated by carbonaceous sputum, cough, hoarseness, or stridor (late). All patients with exposure to smoke or fire in a confined space should receive high-flow oxygen and Pulse Oximetry monitoring.
3. Consider early intubation in patients with respiratory distress, hoarseness, carbonaceous sputum or stridor. If unsure, contact medical command early for assistance with this decision.
4. Confirm and document tube placement with absence of gastric sounds and presence of bilateral breath sounds **AND** confirmatory device (like wave-form ETCO<sub>2</sub> detector). Follow Confirmation of Airway Placement Protocol #2032.
5. For chemical burn exposure, begin flushing immediately with water or appropriate agent for chemical. **Exceptions:** for phosphorous and sodium, **DO NOT** flush with water, cover with cooking oil if available; for Phenol remove with alcohol and follow with large volume of water. If eye is burned, flush with large volume of NSS for 15-20 minutes. May administer tetracaine eye drops before flushing. Continue eye flushing during transport.
6. Monitor ECG for all patients with:
  - a. Electrical/Lightning injury
  - b. Respiratory symptoms
  - c. Multisystem trauma
  - d. Hypovolemic/Traumatic Shock
7. Indicators of severe burn injury include:
  - a. Respiratory tract injury, inhalation injury.
  - b. 2<sup>nd</sup> and 3<sup>rd</sup> degree burns that involve face, hands, feet, genitalia or perineal area or those that involve skin overlying major joints.
  - c. 3<sup>rd</sup> degree burns of greater than 5% BSA.
  - d. 2<sup>nd</sup> degree burns of greater than 15% BSA.
  - e. Significant electrical burns, including lightning injury.
  - f. Significant chemical burns.
  - g. Burn injury in patients with pre-existing illnesses that could complicate management, prolong recovery, or affect mortality.Medical Command physician may direct transport to Burn Center in these cases.
8. **DO NOT** provide fluid bolus if respiratory symptoms are present.
9. Narcotic pain medication should not be given if:
  - a. Oxygen saturation  $\leq$  95%
  - b. SBP  $<$  100 for adults
  - c. SBP  $<$  70 + 2(age in years) for children  $<$  14 y/o
  - d. Patient has altered level of consciousness
10. Reduce dose for patients over 65 y/o.
11. If respiratory depression or hypoxia occur after narcotic:
  - a. Administer oxygen and ventilate if necessary
  - b. If significant respiratory depression, administer naloxone 0.4 mg IV, titrate additional doses until adequate ventilation or total of 2 mg.
12. Nitrous oxide should be self-administered. Patient should be coached to hold mask on his/her face, and the patient will drop mask if he/she becomes sedated. Oversedation may occur if EMS provider holds mask to patient's face.

13. Transport to the closest appropriate medical facility, using the following order of preference:
    - a. If unable to maintain airway or unable to ventilate or patient has symptoms of shortness of breath/cough or inhalation injury is suspected, transport to closest hospital.
    - b. Transport to Trauma Center, if patient has associated trauma. Follow Trauma Destination Protocol #180.
    - c. Transport to a burn center if:
      - 1) The patient has burns to more than 15% BSA or burns to face or hands, and
      - 2) The patient does not meet trauma triage criteria, and
      - 3) The difference between estimated transport time to the closest receiving facility and the burn center is 20 minutes or less.
    - d. If none of the above apply, transport to the closest hospital.
  14. Medical Command Physician may direct transport to Burn Center.
- 

**Performance Parameters:**

- A. Review all burn calls for compliance with Trauma Destinations Protocol # 180
  - B. Review all burn calls for frequency of administration of or documentation of offering pain medication.
-



**THIS PAGE LEFT INTENTIONALLY BLANK**

**HYPOTHERMIA / COLD INJURY / FROSTBITE  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact – Follow Protocol #201  
Assess respirations and pulse for 45 seconds each <sup>1</sup>

- Manage Airway/ Ventilate, as indicated
  - Intubate gently if indicated
- Apply Oxygen <sup>2</sup>
- Monitor ECG / Pulse Oximetry
- Environment
  - Move patient to warm dry place
  - Remove wet clothing
  - Wrap in warm blankets

**Patient temperature > 30° C (86° F)?  
Patient is shivering and conscious?**

**NO**

**YES**

- **TRANSPORT IMMEDIATELY** <sup>3,4</sup>
    - Transport to center capable of bypass rewarming, if possible
    - Consider air ambulance if transport time > 30 minutes
  - Obtain IV/ IO NSS
    - administer NSS 20 mL/kg up to 2000 mL total
    - use warmed NSS if possible
  - Check blood glucose
    - if < 60 mg/dL, administer **Dextrose** <sup>5</sup>
- If cardiac arrest develops, follow Cardiac Arrest-Hypothermia protocol #3035**

- Active external rewarming:
  - apply heat packs to groin, axillae, and neck, if possible. <sup>6</sup>
- Consider IV NSS
- Check blood glucose
  - if < 60 mg/dL, administer **Dextrose**
- If the patient is alert, administer warm non-caffeinated beverages (if available) by mouth slowly. <sup>7</sup>

**Contact Medical Command**

**Contact Medical Command**

Repeat warmed NSS bolus to total of 60 mL/kg or 3000 mL

**HYPOTHERMIA / COLD INJURY / FROSTBITE  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Generalized cooling that significantly reduces the body temperature.
- B. Body temperature < 35° C (95° F).
  - 1. Hypothermia is severe if core body temperature is < 30° C (86° F).
- C. Frostbite generally affects feet, hands, ears, and/or face. Skin initially appears reddened, then mottled, bluish, white and/or gray. This is painful initially then becomes numb.

**Exclusion Criteria:**

- A. Cardiac Arrest from hypothermia – Follow protocol # 3035.
- B. DOA, including the following - see DOA protocol # 322.
  - 1. Submersion for >1 hour.
  - 2. Body tissue/chest wall frozen solid.
  - 3. Body temperature same as surrounding temperature and other signs of death (lividity/ rigor)
- C. Frostbite or cold injury isolated to soft tissues – Follow BLS Hypothermia Protocol # 681

---

**Notes:**

- 1. **Vital signs should be taken for a longer time than usual, so that a very slow pulse or respiratory rate is not missed. Assess pulse for 45 seconds. If a pulse or respirations are detected, do not perform CPR.**
- 2. Use warmed humidified oxygen if available.
- 3. If unresponsive to verbal stimuli or temperature <30° C (86° F), transport to center capable of extracorporeal rewarming (cardiac bypass) if possible. If unsure whether center is capable of 24-hour/7-day emergent bypass rewarming, contact medical command to confirm availability *OR* transport to the closest Level II or III Trauma Center, following Trauma Triage Protocol # 180. Contact medical command at destination facility as soon as possible to provide maximum time for staff to prepare to receive the patient.
- 4. If the patient has severe hypothermia and vertical evacuation is required, transport the patient in a level position when possible. Transporting vertically with the head up has been associated with seizures and death.
- 5. Dextrose dosing:
  - a. Adults- 50% Dextrose, 25 gms IV/ IO
  - b. Pediatrics- 25% Dextrose, 2 mL/kg IV/ IO
- 6. Do not place heat packs directly against skin- wrap in towel.
- 7. **DO NOT** permit fluids by mouth if patient also has severe traumatic injuries or abdominal pain.


---

**Performance Parameters:**

- A. Review for transport to center capable of bypass rewarming when appropriate

**HEAT EMERGENCIES  
STATEWIDE ALS PROTOCOL**

**History/evidence of HEAT exposure**  
**Initial Patient Contact – see Protocol # 201**  
**Check blood glucose and treat hypoglycemia per protocol #7002**  
**Follow Heat Emergency Protocol – see Protocol #686**



**HEAT EMERGENCY  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Heat Cramps** - Painful muscle spasms of the skeletal muscles that occur following heavy work or strenuous exercise in hot environments. Thought to be caused by rapid changes in extracellular fluid osmolarity resulting from fluid and sodium loss. Signs and symptoms include
  - 1. Alert
  - 2. Muscle cramps (normally in muscles most recently heavily exercised)
  - 3. Hot, diaphoretic skin
  - 4. Tachycardia
  - 5. Normotensive
- B. Heat exhaustion** - Patient presents with dizziness, nausea, headache, tachycardia, and possibly syncope. Usually from exposure to high ambient temperatures accompanied by dehydration due to poor fluid intake. Temperature is less than 103° F. Rapid recovery generally follows saline administration.
- C. Heat Stroke**<sup>1</sup> - Patient should be treated as heat stroke if he/she has ALL of the following
  - 1. Exposure to hot environment, and
  - 2. Hot skin, and
  - 3. Altered mental status

**Exclusion Criteria:**

- A. None

**Possible MC Orders:**

- A. Medical command physician may order release of care for mild heat cramps or mild heat exhaustion.
- B. May order additional fluid boluses


---

**Notes:**

- 1. Patient's thermoregulatory mechanisms break down completely. Body temperature is elevated to extreme levels, which results in multi-system tissue damage including altered mental status. Heat stroke often affects elderly patients with underlying medical disorders. Patients usually have dry skin; however, up to 50% of patients with exertional heat stroke may exhibit persistent sweating. Therefore, patients with heat stroke may be sweating.
  - 2. Patient may take oral fluid replacement rather than IV if no nausea. Allow oral intake of cool fluids or water (may use commercial sport/rehydration drinks like Gatorade or Powerade) if patient is alert. Do not permit the patient to drink if altered mental status, abdominal pain or nausea. Avoid carbonated sodas, alcoholic beverages, and caffeinated beverages.
-

### ALTERED LEVEL OF CONSCIOUSNESS STATEWIDE ALS PROTOCOL

Initial Patient Contact - See Protocol # 201  
Administer Oxygen <sup>1</sup>  
Manage Airway/Ventilate, if needed <sup>2,3</sup>  
Monitor ECG/Pulse Oximetry <sup>4</sup>  
Assess Glasgow Coma Scale  
Initiate IV/IO NSS  
Draw blood <sup>5</sup>



**ALTERED LEVEL OF CONSCIOUSNESS - Adult  
STATEWIDE ALS PROTOCOL****Criteria:**

- A.** Patient with altered level of consciousness due to:
1. Unclear etiology after assessing patient
  2. History consistent with hypoglycemia
  3. Suspected drug ingestion /overdose

**Exclusion Criteria:**

- A.** Altered level of consciousness due to:
1. Trauma - Follow appropriate trauma protocol (e.g. head injury or multi-system trauma protocol)
  2. Shock - Follow Shock protocol # 7005
  3. Dysrhythmias - Follow appropriate dysrhythmia protocol.
  4. Toxicologic
    - a. Carbon monoxide - Follow Poisoning/Toxic Exposure Protocol #8031.
    - b. Cyanide - Follow Cyanide Exposure Protocol #8081.
    - c. Nerve agent exposure - Follow Nerve Agent Exposure Protocol #8083.
  5. Seizure - Follow Seizure Protocol #7007.
  6. Stroke - Follow Stroke Protocol #7006.
  7. Other medical problems specifically suspected due to history or exam, e.g. choking, hypoxia due to respiratory failure, etc...- Follow applicable specific protocol.

**Possible MC Orders:**

- A.** Additional doses of naloxone
- B.** Additional doses of dextrose or glucagon (if available)

**Notes:**

1. Administer oxygen by appropriate method.
2. Confirm and document tube placement with auscultation and ETCO<sub>2</sub> detector/secondary device - Follow Confirmation of Airway Placement Protocol #2032
3. If unable to intubate on up to 3 attempts, consider Alternative/ Rescue airway.
4. See Pulse Oximetry Protocol #226. Pulse Oximetry must not delay the application of oxygen. Record SpO<sub>2</sub> after administration of oxygen or intubation.
5. If receiving facility will accept blood samples, blood should be drawn in red top tube for analysis at the hospital unless the patient is a known diabetic who takes insulin or oral diabetic medications (e.g. micronase, glyburide, glucophage, etc...)
6. Indications of possible opiate overdose include decreased respirations, pinpoint pupils, skin "track marks", *AND/OR* the presence of drug paraphernalia.
7. Naloxone should not be given to patients that have been intubated.
8. Naloxone can be administered IM, IO, or intranasally if IV cannot be established. Ideally, intranasal administration should be done via an atomizing device. If IM route is required, use 2 mg.
9. Dose should be titrated to improved respirations. Larger individual doses of naloxone can precipitate opiate withdrawal with the potential for a violent or combative patient that is difficult to manage at the scene and once the patient is admitted to the hospital. If no response to dose of naloxone, dose may be repeated in 0.4 mg increments to a total of 2 mg. Some opioids, like propoxyphene and dextromethorphan, require higher doses of naloxone.
10. Indicators of improved mental status include:
  - a. Orientation to person, place and time
  - b. Increased alertness
  - c. Increased responsiveness to questions
11. For patients refusing transport, adhere to Refusal of Treatment /Transport Protocol #111.
12. Patient may be released without Medical Command if all of the following are met in addition to criteria in protocol #111:
  - a. Repeat glucose meter is > 80 mg/dl
  - b. Patient is an insulin-dependent diabetic (not on oral antihyperglycemics)
  - c. Patient returns to normal mental status, with no focal neurologic signs/symptoms after receiving glucose

- d. Patient can promptly obtain and will eat a carbohydrate meal.
  - e. Patient refuses transport, or patient and paramedics agree transport not needed
  - f. Another competent adult will be staying with patient
  
  - g. No major co-morbid conditions exist, such as chest pain, arrhythmias, dyspnea, seizures, intoxication
  - h. Patient should not be released without medical command contact if given glucagon instead of dextrose or if he/she received naloxone.
  - i. If all of the above conditions are not met and the patient or legal guardian refuses transport, contact medical command. If the patient or legal guardian requests transport, honor the request.
- 

**Performance Parameters:**


- A. Review for proper use of naloxone and glucose and documentation of neurologic assessment/response to treatment.
-



**THIS PAGE LEFT INTENTIONALLY BLANK**

**ALTERED LEVEL OF CONSCIOUSNESS - Pediatric  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - See Protocol #201  
Administer Oxygen <sup>1</sup>  
Manage Airway/Ventilate, if needed <sup>2,3</sup>  
Monitor ECG/Pulse Oximetry <sup>4</sup>  
Assess Glasgow Coma Scale  
Initiate IV/IO NSS  
Draw blood <sup>5</sup>



**ALTERED LEVEL OF CONSCIOUSNESS - Pediatric  
STATEWIDE ALS PROTOCOL****Criteria:**

- A.** Patient with altered level of consciousness due to:
1. Unclear etiology after assessing patient
  2. History consistent with hypoglycemia (in infants and children, hypoglycemia frequently accompanies overdose, alcohol ingestion, poisoning, or metabolic/medical diseases)
  3. Suspected drug ingestion /overdose

**Exclusion Criteria:**

- A.** Altered level of consciousness due to:
1. Trauma - Follow appropriate trauma protocol (e.g. head injury or multi-system trauma protocol)
  2. Shock - Follow Shock Protocol #7005
  3. Dysrhythmias - Follow appropriate dysrhythmia protocol.
  4. Toxicologic
    - a. Carbon monoxide - Follow Poisoning / Toxic Exposure Protocol #8031.
    - b. Cyanide - Follow Cyanide Exposure Protocol #8081.
    - c. Nerve agent exposure - Follow Nerve Agent Exposure Protocol #8083.
  5. Seizure - Follow Seizure Protocol #7007.
  6. Stroke - Follow Stroke Protocol #7006.
  7. Other medical problems specifically suspected due to history or exam, e.g. choking, hypoxia due to respiratory failure, etc...- Follow applicable specific protocol.

**Possible MC Orders:**

- A.** Additional doses of naloxone
- B.** Additional doses of dextrose or glucagon (if available)

---

**Notes:**

1. Administer oxygen by appropriate method.
2. In children, ventilation by bag-valve-mask is the preferred method of airway maintenance and ventilation if transport time is short. However, if patient cannot be adequately oxygenated or ventilated by bag-valve-mask or if transport time is long, intubation is indicated. Use a length-based device to assist with selection of appropriate sized airway equipment.
3. Confirm and document tube placement with auscultation and ETCO<sub>2</sub> detector/secondary device - Follow Confirmation of Airway Placement Protocol #2032
4. See Pulse Oximetry Protocol #226. Pulse Oximetry must not delay the application of oxygen. Record SpO<sub>2</sub> after administration of oxygen or intubation.
5. Blood should be drawn in red top tube for analysis at the hospital unless the patient is a known diabetic who takes insulin or oral diabetic medications (e.g. micronase, glyburide, glucophage, etc...)
6. Indications of possible opiate overdose include decreased respirations, pinpoint pupils, skin "track marks", *AND/OR* the presence of drug paraphernalia.
7. Naloxone should not be given to patients that have been intubated.
8. Naloxone can be administered IM, IO, or intranasally if IV cannot be established. Ideally, intranasal administration should be done via an atomizing device.
9. Dose should be titrated to improved respirations. Larger individual doses of naloxone can precipitate opiate withdrawal with the potential for a violent or combative patient that is difficult to manage at the scene and once the patient is admitted to the hospital. If no response to dose of naloxone, dose may be repeated in 0.4 mg increments to a total of 2 mg. Some opioids, like propoxyphene and dexamethorphan, require higher doses of naloxone.
10. Indicators of improved mental status include:
  - a. Orientation to person, place and time
  - b. Increased alertness
  - c. Increased
  - d. If no response to dose responsiveness to questions
11. For patients refusing transport, adhere to Refusal of Treatment/Transport Protocol #111.
12. For neonates, 25% dextrose dose should be diluted with equal amounts of NSS for 12.5% dextrose at 4 mL/kg.

13. Patient may be released without Medical Command if all of the following are met in addition to criteria in protocol #111:
  - a. Repeat glucose meter is > 60 mg/dl
  - b. Patient is an insulin-dependent diabetic (not on oral antihyperglycemics)
  - c. Patient returns to normal mental status, with no focal neurologic signs/symptoms after receiving glucose.
  - d. Patient can obtain and will promptly eat a carbohydrate meal.
  - e. Legal guardian refuses transport, or patient, legal guardian and paramedics agree transport not needed
  - f. Legal guardian or another competent adult will be staying with patient
  - g. No major co-morbid conditions exist, such as chest pain, arrhythmias, dyspnea, seizures, intoxication
  - h. Patient should not be released without medical command contact if given glucagon instead of dextrose or if he/she received naloxone.
  - i. If all of the above conditions are not met and the patient or legal guardian refuses transport, contact medical command. If the patient or legal guardian requests transport, honor the request.

---


**Performance Parameters:**

- A. Review for proper use of naloxone and glucose and documentation of neurologic assessment/response to treatment.

**THIS PAGE LEFT INTENTIONALLY BLANK**

**SHOCK  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact - Follow protocol #201  
 Manage Airway/Ventilate, if needed <sup>1,2,3</sup>  
 High-flow oxygen  
 Keep patient warm  
 Monitor ECG/Pulse Oximetry <sup>4</sup>



## SHOCK STATEWIDE ALS PROTOCOL

**Criteria:**

- A. Hypoperfusion of body organs is characterized by alterations in mental status, pallor, diaphoresis, tachypnea, tachycardia, poor capillary refill, and hypotension.

For example:

1. Septic Shock - signs or symptoms of hypoperfusion from a suspected infectious source (e.g. urosepsis, pneumonia, bacteremia / septicemia). These patients may present with a fever or preceding infectious illness.
2. Hypovolemic Shock from gastrointestinal bleeding or from repetitive vomiting/diarrhea in infants/children.
3. Hypoperfusion of unknown etiology.

**Exclusion Criteria:**

- A. Cardiogenic Shock- hypotension with suspected pulmonary edema - See CHF Protocol #5002.
- B. Hypovolemic/Traumatic Shock of traumatic etiology - See Multisystem Trauma or Traumatic Shock Protocol #6002.
- C. Neurogenic Shock due to spinal cord injury – See Multisystem Trauma or Traumatic Shock Protocol #6002.

**Possible MC Orders:**

- A. Additional NSS fluid boluses
- B. Earlier intervention with vasopressor infusions (dopamine, dobutamine, epinephrine).



---

**Notes:**

1. Confirm and document tube placement with auscultation and ETCO<sub>2</sub> detector/secondary device - Follow Confirmation of Airway Placement Protocol #2032
  2. If unable to intubate on up to 3 attempts, consider alternative/ rescue airway device.
  3. In children, ventilation by bag-valve-mask is the preferred method of airway maintenance and ventilation if transport time is short. However, if patient cannot be adequately oxygenated or ventilated by bag-valve-mask or if transport time is long, intubation is indicated. Use a length-based device to assist with selection of appropriate sized airway equipment.
  4. See Pulse Oximetry Protocol #226. Pulse Oximetry must not delay the application of oxygen. Record SpO<sub>2</sub> after administration of oxygen or intubation.
  5. Bolus IV fluid should be given as quickly as possible, ideally in less than ten minutes.
  6. Do not give IV fluid bolus prior to medical command if the patient has signs of CHF (for example, rales or significant pitting edema).
  7. Some recommendations suggest using dobutamine for mild cardiogenic shock (SBP 70-90) and dopamine for severe shock (SBP < 70). Mix infusion using regional or service prescribed concentration, and administer 5-20 mcg/kg/min. Generally start at 5 mcg/kg/min, and increase every 10 minutes by an additional 5 mcg/kg/min until SBP >100 mmHg. **DO NOT exceed 20 mcg/kg/min unless ordered by medical command physician.**
  8. If unable to obtain peripheral IV access, place an intraosseous (IO) line, if available.
  9. In infants, it is difficult to distinguish between hypoperfusion from hypovolemia and that due to cardiogenic shock. Hypovolemia frequently follows a history of repetitive vomiting/diarrhea. If cardiogenic shock is suspected, fluid boluses should be limited to the initial 20 mL/kg.
-

**STROKE  
STATEWIDE ALS PROTOCOL**



Initial Patient Contact - See Protocol #201  
Administer Oxygen<sup>1</sup>  
Manage Airway/Ventilate, if needed<sup>2,3</sup>  
Monitor ECG/Pulse Oximetry<sup>4</sup>



**Is acute stroke suspected by Cincinnati Prehospital Stroke Scale<sup>5,6</sup> (CPSS)?**

Face - facial droop present,  
*OR*  
Arm - upper extremity arm drift present (arms extended/ palms up),  
*OR*  
Speech - inability to say, "The sky is blue in Pennsylvania" normally,  
*AND*  
Time - time of symptom onset definitely < 3 hours<sup>7</sup>

Exclude patient if another history of a stroke within last 3 months,  
*OR*  
Major surgery within last 14 days.





## STROKE STATEWIDE ALS PROTOCOL

**Criteria:**

- A.** Patients may have the following clinical symptom(s):
1. Impaired expression or understanding of speech
  2. Unilateral weakness/hemiparesis
  3. Facial asymmetry/droop
  4. Headache
  5. Poor coordination or balance
  6. Partial loss of peripheral vision
  7. Vertigo
- B. CAUTION:** Respiratory and cardiovascular abnormalities may reflect increased intracranial pressure. Lowering of the blood pressure may be dangerous.

**Exclusion Criteria:**

- A.** Consider hypoglycemia, trauma, and other etiologies that can cause focal neurological symptoms that mimic stroke, and follow applicable protocol if appropriate.

**Possible MC Orders:**

- A.** Transport to a receiving hospital that is not the usual or closest hospital because the medical command physician has knowledge that another facility is able to better treat an acute stroke within a critical time window.

---

**Notes:**

1. Administer oxygen by appropriate method and monitor Pulse Oximetry, if available.
2. Confirm and document tube placement with auscultation and ETCO<sub>2</sub> detector/secondary device - Follow Confirmation of Airway Placement Protocol #2032
3. If unable to intubate on up to 3 attempts, consider alternative/ rescue airway device.
4. See Pulse Oximetry Protocol #226. Pulse Oximetry must not delay the application of oxygen. Record SpO<sub>2</sub> after administration of oxygen or intubation.
5. Neurological examination includes level of consciousness, Glasgow Coma Scale, pupils, individual limb movements, and Cincinnati Prehospital Stroke Scale (CPSS).
6. **Cincinnati Prehospital Stroke Scale.** If any of the following is **abnormal** and **new** for the patient, he/she may have an acute stroke:
  - a. Facial Droop (patient smiles or shows teeth) - abnormal if one side of the face does not move as well as the other.
  - b. Arm Drift (patient holds arms straight out in front of him/her and closes eyes) – abnormal if one arm drifts down compared with the other.
  - c. Speech (patient attempts to say “The sky is blue in Pennsylvania”) – abnormal if patient slurs words, uses inappropriate words, or can’t speak.
7. Attempt to identify the precise time of the onset of the patient’s first symptoms. The time of onset is extremely important information, and patient care may be different if patient can be delivered to a receiving hospital capable of treating acute strokes within 3 hours from onset of symptoms. Time is based upon the last time that the patient was witnessed to be at his/her neurologic baseline.
8. Transport and Medical Command contact should not be delayed by attempts to initiate IV or draw blood in patients who are awake. In these patients, the IV should be done enroute after notifying receiving facility or medical command.
9. If patient can be delivered by air (but not by ground) to receiving facility within 3 hours of symptom onset, consider contact with medical command for assistance in deciding upon the utility of air medical transport.
10. If patient can’t tolerate supine position, transport with head elevated < 30 degrees.
11. Contact Medical Command for all patients with acute CPSS symptoms that have onset within 3 hours of estimated arrival at the receiving facility, so the receiving hospital can prepare for the patient’s arrival. Describe to the Medical Command Physician your findings, including CPSS results. Medical command may order transport to a facility other than the closest facility if another center is better prepared to evaluate and treat an acute stroke. If the medical command physician is not at the receiving facility, the medical command physician should relay pertinent information to the receiving facility.
12. If patient will arrive for ED treatment within 3 hours of symptoms, initiate a second IV access with saline lock enroute to hospital. Ideally 18-20 gauge IV access.

13. Before administering glucose, blood should be drawn in red top tube for analysis at the hospital unless the patient is a known diabetic who takes insulin or oral diabetic medications (e.g. micronase, glyburide, glucophage, etc...).
  14. If Glucometer < 60 mg/dL, give 50% dextrose 25 gm. IV.
- 

**Performance Parameters:**

- A. Review on scene time for all cases of suspected stroke with time of symptom onset less than 3 hours from time of EMS arrival. Consider benchmark of on scene time  $\leq 10$  minutes.
  - B. Review documentation for CPSS criteria, time of symptom onset, glucose determination, and appropriate communication with medical command and receiving facility to maximize prearrival warning to receiving facility and most appropriate receiving facility.
-

**THIS PAGE INTENTIONALLY LEFT BLANK**

**SEIZURE  
STATEWIDE ALS PROTOCOL**

**Initial Patient Contact - See Protocol #201<sup>1</sup>**  
**If history/evidence of trauma, maintain c-spine stabilization**


**(Follow C-Spine Immobilization Protocol if indicated)**

**Administer Oxygen<sup>2</sup>**

**Manage Airway/Ventilate, if needed<sup>3,4</sup>**

**Monitor ECG/Pulse Oximetry, if seizure permits<sup>5</sup>**

**Initiate IV/IO NSS, if possible**



## SEIZURE STATEWIDE ALS PROTOCOL

**Criteria:**

- A. Patients who are actively seizing with generalized clonic-tonic seizure. Indicators of seizures requiring treatment include:
  - 1. two or more consecutive seizures without return of consciousness between episodes.
  - 2. ongoing seizure for more than 4 minutes.
  - 3. seizures associated with hypoxia.
- B. Patients who have had tonic-clonic seizure activity prior to EMS arrival.

**Exclusion Criteria:**

- A. Patients who are postictal following a single seizure and have history or evidence of trauma - Follow Multi-system Trauma or Traumatic Shock Protocol #6002 or Head Injury Protocol #611, as indicated.

**Possible MC Orders:**

- A. May order additional doses of benzodiazepine.
- B. May order lidocaine.

---

**Notes:**

1. Determine (if possible):
  - a. Type of seizure:
    - Generalized
    - Focal
  - b. Stage of seizure:
    - Active
    - Postictal
  - c. Cause of seizure:
    - Infections
    - Drug overdose
    - Metabolic
    - Hypoxia
    - Toxins
    - Stroke
    - Traumatic
    - Vascular
    - Alcohol withdrawal
    - Non-compliance with medications
2. Administer oxygen by appropriate method and monitor Pulse Oximetry, if available. Patients with ongoing seizure activity should receive high-flow oxygen.
3. Confirm and document tube placement with auscultation and ETCO<sub>2</sub> detector/secondary device - Follow Confirmation of Airway Placement Protocol #2032
4. If unable to intubate on up to 3 attempts, consider alternative/ rescue airway.
5. See Pulse Oximetry protocol #226. Pulse Oximetry must not delay the application of oxygen. Record SpO<sub>2</sub> after administration of oxygen or intubation.
6. Blood should be drawn in red top tube for analysis at the hospital unless the patient is a known diabetic who takes insulin or oral diabetic medications (e.g. micronase, glyburide, glucophage, etc...), has a known history of seizure disorder, or has ongoing seizure activity that prohibits blood draw.
7. Prevent patient from sustaining physical injury.
8. 50% Dextrose may be diluted 1:1 with NSS to administer 25% Dextrose
9. Glucagon dosage (if available):
  - a. 1 mg IM if patient is  $\geq$  20 kg or 5 y/o
  - b. 0.5 mg IM if patient is  $<$  20 kg or 5 y/o
10. Seizures related to eclampsia can occur in the third trimester or can even occur days or weeks after delivery. Eclampsia should be considered in pregnant or post-partum women who have a new onset seizure without prior history of seizure disorder or who have a history of preeclampsia or hypertension associated with the pregnancy.
11. If eclampsia seizure does not stop after magnesium, then administer benzodiazepine as listed.
12. If IV/ IO is not obtainable, may administer rectal or IM medications. May repeat these doses once.


---

**Performance Parameters:**

- A. Review for documentation of blood glucose if patient does not have a history of seizure disorder.
- B. Review for documentation of vital signs and Pulse Oximetry after administration of benzodiazepine.
- C. Review for documentation of description of any witnessed seizure activity.

**SERIOUSLY ILL APPEARING PATIENT  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact- See Protocol #201  
Initiate IV NSS  
If signs of hypoglycemia, check blood glucose  
Consider obtaining blood samples



**SERIOUSLY ILL APPEARING PATIENT  
STATEWIDE ALS PROTOCOL****Criteria:**

- A.** Any situation not covered under another existing protocol, in which the practitioner determines that the patient is potentially seriously ill with a condition that may suddenly deteriorate with the possibility of requiring the administration of medications or fluids.

**Exclusion Criteria:**

- A.** Patient is stable and no ALS intervention is anticipated.
- 

**Performance Parameters:**

- A.** Review for stable patients with no indication for necessity of initiating IV access.
- B.** Review for specific documentation of need for IV.

**NAUSEA / VOMITING  
STATEWIDE ALS PROTOCOL**

Initial Patient Contact- See Protocol #201


Initiate IV NSS

Administer NSS bolus of 20 mL /kg (2000 mL max)

**(unless contraindicated by h/o CHF, renal failure, etc)**

Check blood glucose

Consider drawing blood samples



**Benzodiazepine Options <sup>3</sup>:**  
**(ADULTS ONLY, choose one)**  
 Midazolam 1 mg IV/ IO slowly  
*OR*  
 Diazepam 5 mg IV/ IO slowly  
*OR*  
 Lorazepam 1mg IV/ IO slowly



**NAUSEA / VOMITING  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Patient with persistent nausea or vomiting.

**Exclusion Criteria:**

- A. Patient is stable and no ALS intervention is anticipated.
- B. Hypotension- See Shock Protocol # 7005.
- C. Altered mental status – See Altered Level of Consciousness Protocol #7002A or #7002P.

**Possible Medical Command Orders:**

- A. For children between 6 m/o –14 y/o, may order ondansetron (if available) 0.1 mg/kg IM/IV (maximum dose of 4 mg).
- 

**Performance Parameters:**

- A. Review for contact with Medical Command before giving ondansetron to patients who are < 14 y/o.

**POST-PARTUM HEMORRHAGE  
STATEWIDE ALS PROTOCOL**

See Emergency Childbirth Protocol # 781  
Assure all fetuses have been delivered

Administer Oxygen  
Firmly massage the uterus

Initiate IV/IO NSS, 500 mL bolus  
(If hypotension, administer up to 2000 mL NSS at wide-open rate)  
Monitor ECG/Pulse Oximetry

Contact Medical  
Command

Oxytocin IV infusion (if available)  
10-20 units/1000mL NSS at wide open rate

If hypotension  
Follow Shock Protocol #7005

**POST-PARTUM HEMORRHAGE  
STATEWIDE ALS PROTOCOL**

**Criteria:**

- A. Excessive uterine bleeding after delivery of neonate (continued steady flow of bright red blood)
- B. Uterine bleeding and signs of hypoperfusion after delivery of neonate

**Exclusion Criteria:**


- A. Patient known to be pregnant with multiple fetuses (more than delivered)
- B. Patient who has not had a prenatal ultrasound to confirm the number of fetuses.

**Possible MC Orders:**

- A. Oxytocin IV infusion (if available), 10-20 units / 1000 ml NSS at wide-open rate.

**AGITATED BEHAVIOR / PSYCHIATRIC DISORDERS  
STATEWIDE ALS PROTOCOL**

Follow BLS Agitated Behavior/Psychiatric Disorders Protocol #801  
 The safety of EMS personnel is top priority  
 Wait for law enforcement if scene is unsafe  
 Request Law Enforcement as needed



**Sedation Options:**  
(Choose one)

**Lorazepam** 1-2 mg IM/IV/IO <sup>3</sup> (0.1 mg/kg, max 2 mg/dose)  
may repeat every 5 minutes until maximum of 4 mg

*OR*

**Diazepam** 5-10 mg IM/IV/IO <sup>3</sup> (0.1 mg/kg)  
may repeat every 5 minutes until maximum 0.3 mg/kg

*OR*

**Midazolam** 1-5 mg IM/IV/IO <sup>3,4</sup> (0.05 mg/kg)  
may repeat every 5 minutes until maximum of 0.1 mg/kg

**AGITATED BEHAVIOR / PSYCHIATRIC DISORDERS  
STATEWIDE ALS PROTOCOL****CRITERIA:**

- A.** Patient with a psychiatric or behavioral disorder who is at imminent risk of self-injury or is a threat to others.

*OR*

- B.** Patient with a medical condition causing agitation and possibly violent behavior. Examples of these conditions are:
1. Alcohol or drug (e.g. PCP, methamphetamine, cocaine) intoxications
  2. Hypoglycemia
  3. Stroke
  4. Drug overdose
  5. Post-ictal after seizure
  6. Head trauma

**Procedure for patients that require physical restraint:****A. All Patients:**

1. Use the minimum amount of restraint necessary to safely accomplish patient care and transportation with regard to the patient's dignity.
2. Assure that adequate personnel are present and that police assistance has arrived, if available, before attempts to restrain patient.
3. Restrain all 4 extremities with patient supine on stretcher.<sup>5,6,7,8</sup>
4. Use soft restraints to prevent the patient from injuring him or herself or others.<sup>9</sup>
  - a. If the handcuffs or law enforcement devices are used to restrain the patient, a law enforcement officer should accompany the patient in the ambulance
  - b. It is preferable that a law enforcement officer follows the ambulance in a patrol car to the receiving facility if physical restraint is necessary.
5. Do not place restraints in a manner that may interfere with evaluation and treatment of the patient or in any way that may compromise patient's respiratory effort.<sup>10</sup>
6. If the patient is spitting, may cover his/her face with a surgical mask or with a NRB mask with high flow oxygen.<sup>11</sup>
7. Evaluate circulation to the extremities frequently.
8. Thoroughly document reasons for restraining the patient, the restraint method used, and results of frequent reassessment.

**Possible Medical Command Orders:**

- A.** Additional benzodiazepine

---

**Notes:**

1. Verbal techniques include:
  - a. Direct empathetic and calm voice.
  - b. Present clear limits and options.
  - c. Respect personal space.
  - d. Avoid direct eye contact.
  - e. Non-confrontational posture.

2. Do not permit patient to continue to struggle against restraints. This can lead to death due to severe rhabdomyolysis, acidosis, dysrhythmia, or respiratory failure. Medical command should be contacted for possible chemical restraint with sedative medication.
3. If age > 65, reduce doses of sedative benzodiazepines in half.
4. Regional or service policy may permit intranasal midazolam, but this may not be as effective as parenteral medications.
5. Initial “take down” may be done in a prone position to decrease the patient’s visual field and ability to bite, punch, and kick. After the individual is controlled, he/she should be restrained to the stretcher or other transport device in the supine position.
6. **DO NOT restrain patient in a hog-tied or prone position.**
7. **DO NOT** sandwich patient between devices, such as long boards or Reeve’s stretchers, for transport. Avoid restraint to unpadded devices like backboards.
8. A stretcher strap that fits snugly just above the knees is effective in decreasing the patient’s ability to kick.
9. Padded or leather wrist or ankle straps are appropriate. Handcuffs and plastic ties are not considered soft restraints.
10. Never apply restraints near the patient’s neck or apply restraints or pressure in a fashion that restricts the patient’s respiratory effort.
11. Never cover a patient’s mouth or nose except with a surgical mask or a NRB mask with high flow oxygen. A NRB mask with high flow oxygen may be used to prevent spitting in a patient that also may have hypoxia or another medical condition causing his/her agitation, but a NRB mask should never be used to prevent spitting without also administering high flow oxygen through the mask.


---

**Performance Parameters:**

- A. Review every case of the use of physical or chemical restraint for documentation of physical restraint procedure, monitoring of respiratory effort, assessment of extremity neurovascular status every 15 minutes, and medical command physician orders for use of physical or chemical restraint.



**THIS PAGE INTENTIONALLY LEFT BLANK**

**POISONING/TOXIN EXPOSURE  
STATEWIDE ALS PROTOCOL**







**POISONING/TOXIN EXPOSURE  
STATEWIDE ALS PROTOCOL**



**POISONING/TOXIN EXPOSURE  
STATEWIDE ALS PROTOCOL**



**POISONING/TOXIN EXPOSURE  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patient who has accidentally or purposefully been exposed to toxic substances. Including:
1. **Ingested toxins**
    - a. For example pills, capsules, medications, recreational drugs, poisonous plants, strong acids or alkali household or industrial compounds.
  2. **Inhaled toxins**
    - a. For example carbon monoxide and other toxic gases.
  3. **Absorbed toxins**
    - a. For example substances on skin or splashed into eyes.
  4. **Injected toxins**
    - a. For example snake bites or substances injected through the skin.

**Exclusion Criteria:**

- A. Patient with altered level of consciousness (unless suspected carbon monoxide poisoning) - follow Altered Level of Consciousness Protocol #7002A or #7002P.
- B. Patient with exposure to organophosphate pesticide or nerve agent – follow Nerve Agent Exposure Protocol #8083.
- C. Patient with exposure to cyanide – follow Cyanide Exposure Protocol #8081.
- D. Patient with suspected allergic reaction/anaphylaxis – follow Allergic Reaction Protocol #4011.

**Treatment:****A. All patients:**

1. Initial Patient Contact – see Protocol #201.
  - a. **WARNING: EMS personnel must not enter confined spaces with potential toxic gases (e.g. manure pits, silos, spaces with carbon monoxide, spaces with industrial gases) unless personnel have proper training and PPE.**
  - b. **Decontamination- Ideally, patients will be fully decontaminated before treatment and transport. In situations where the patient has not been fully decontaminated, EMS personnel must be in appropriate PPE before treatment/ transport. The receiving ED should be notified ASAP so that they can prepare for any additional decontamination that is needed on arrival.**
  - c. If toxic exposure/overdose is the result of intentional behavior - also see Agitated Behavior/Psychiatric Disorders protocol # 8001. <sup>1</sup>
2. Maintain adequate airway.
3. Administer oxygen by appropriate method and monitor Pulse Oximetry <sup>2</sup>, if available.
  - a. (High-flow oxygen if suspected carbon monoxide poisoning, respiratory distress, or cough).
4. Determine:
  - a. **What** – identify specific toxin and amount, if possible.
    - 1) If possible, safely transport source of toxin (e.g. prescription pill bottles) with patient to receiving facility.
    - 2) EMS services should not transport dangerous items (e.g. toxic chemicals that are not sealed in their original containers, live snakes, etc...)
  - b. **When** – identify time of exposure, if possible.
  - c. **Why** – identify reason for exposure, if possible.
  - d. **Where** – identify environmental site issues (e.g. exposure in a confined space or carbon monoxide present).
5. Treat specific toxins based upon the appropriate category:
  - a. **Ingested Toxins.** Treat all exposures as follows:
    - 1) **DO NOT INDUCE VOMITING.**
    - 2) For asymptomatic ingestions that may not require transport, consider contacting Poison Control Center. <sup>3,4</sup>
    - 3) Initiate IV/IO NSS KVO if patient has symptoms.
      - a) If hypotensive, follow Shock Protocol #7005.
    - 4) If mental status changes, then check blood glucose and treat hypoglycemia per Altered Level of Consciousness Protocol #7002A or #7002P.
    - 5) Monitor ECG

- 6) Contact Medical Command for possible order for activated charcoal.<sup>3,4,5</sup>
- b. **Inhaled Toxins.** Treat all symptomatic (e.g. SOB, cough, headache, decreased LOC) patients as follows:
  - 1) Only personnel with proper training and wearing proper PPE should enter environments that may have toxic gases.
  - 2) Remove patient from environment.
  - 3) Administer 100% oxygen.
  - 4) Ventilate with BVM, if needed.
  - 5) Intubate if indicated.
  - 6) Initiate IV/IO NSS KVO
    - a) If hypotensive, follow Shock Protocol # 7005
  - 7) Monitor ECG and Pulse Oximetry
    - a) **WARNING: Pulse Oximetry monitors give false readings in patients that have been exposed to carbon monoxide or cyanide, and normal readings should not diminish the use of 100% oxygen in these patients.**
  - 8) Consider transport to a facility with hyperbaric oxygen if patient has suspected carbon monoxide exposure and has altered mental status or is pregnant. This may be appropriate if time to transport to a facility with hyperbaric oxygen capability is within 30 minutes of the time to transport to the closest facility and the patient is deemed to be stable for the longer transport.
  - 9) Contact Medical Command
- c. **For Absorbed Toxins:**
  - 1) Remove contaminated clothing.
  - 2) Flush affected area copiously:
    - a) Liquid substance - Irrigate with copious amounts of room temperature water. Do not contaminate uninjured areas while flushing.
    - b) Dry substances - With gloves and appropriate PPE, brush remaining powder from skin and clothing, then irrigate with copious amounts of water.<sup>6</sup>
    - c) **Eyes** - Flush affected eyes continuously with water or saline if eye exposure. **Prior to flushing eyes, may instill 1-2 drops of ophthalmic tetracaine (if available) and may repeat every 15 minutes as needed for comfort. Do not instill tetracaine or irrigate eyes if there has been blunt trauma/ possible penetrating injury to the eye.**
- d. **For Injected Poisons/Snakebite:**
  - 1) Identify type of snake or animal (e.g. scorpion), if safe and possible. If identity of a snake is not known, all victims of snakebite should be treated as if the snake is poisonous. Do not delay transport or endanger individuals by attempting to capture or kill a snake.
  - 2) Calm patient.
  - 3) Administer high-flow oxygen, if respiratory symptoms are present.
  - 4) Remove jewelry and tight clothing.
  - 5) Consider immobilizing the involved body part. If extremity involved, keep the extremity below the level of the patient's heart.
  - 6) Keep the patient as still as possible to reduce the circulation of the venom. Carry patient for transport, if possible.
  - 7) Apply constricting band proximal to bite if patient is hypotensive. **DO NOT APPLY TOURNIQUET.**
  - 8) **DO NOT APPLY ICE.**
  - 9) Initiate IV/IO NSS KVO and draw blood (including tubes for type and cross, if available)
    - a) If hypotensive, follow Shock Protocol - #7005
  - 10) Contact Medical Command.

**Possible Medical Command Orders:**

- A. Administration of activated charcoal, **if available**, may be ordered <sup>4,5</sup>:
  1. **Adults:** 25 - 50 gm orally of pre-mixed activated charcoal.
  2. **Children:** 1 gm/kg orally or approximately 12.5 - 25 gm orally of pre-mixed activated charcoal.
- B. If tricyclic antidepressant overdose and patient hypotensive, may order sodium bicarbonate.
- C. If calcium channel blocker or beta-blocker overdose and hypotensive, may order calcium chloride (if available) or glucagon (if available).

- D. If dystonic reaction, may order diphenhydramine.
  - E. If smoke inhalation (cyanide risk) or suspected asphyxiation from hydrogen sulfide (e.g. in manure pit), may order sodium thiosulfate (if available).
  - F. If suspected carbon monoxide toxicity and altered level of consciousness or pregnant, may order transport to center capable of hyperbaric oxygen therapy.
- 

**Notes:**

1. Patients who have ingested a toxic substance with suicidal intent may not refuse transport. See Refusal of Treatment/Transport Protocol #111.
  2. See Pulse Oximetry Protocol #226. Pulse Oximetry is not accurate in patients with suspected exposure to carbon monoxide or cyanide and shall not be used in these situations.
  3. **National Poison Control Center telephone number is 800-222-1222.** EMS personnel must follow instructions from Poison Control Center unless the orders are superceded by orders from a medical command physician. These instructions must be documented on the PaPCR. Poison Control Center should only be contacted for stable patients with minor ingestions. Medical Command should be contacted for patients who are likely to require transportation to a hospital.
  4. Activated charcoal may only be given by order of medical command or poison control.
  5. Contraindications to charcoal:
    - a. Patient unable to swallow/protect airway.
    - b. Seizures.
    - c. Hydrocarbons ingestion (e.g. turpentine)
    - d. Caustic substance ingestion (e.g. liquid drain cleaner or milk pipe cleaner)
  6. **Note** - some substances, like dry lime will cause a heat-producing reaction when mixed with water. Copious water should be available before beginning to irrigate.
- 

**Performance Parameters:**

- A. Review for documentation of orders received from Poison Control Centers or Medical Command.

### CYANIDE COMPOUND EXPOSURE STATEWIDE ALS PROTOCOL

#### Decontaminate patient

If possible, treat patients with severe exposure during decontamination

**CAUTION: Only personnel wearing Level B PPE (with appropriate training) should treat patient before decontamination**

Initial Patient Contact - See protocol #201  
Manage Airway/Ventilate, if needed <sup>1</sup>  
Administer High-flow Oxygen <sup>1</sup>

**NOTE: Pulse Oximetry may be inaccurate and should be avoided**

**Patient with decreased LOC, seizures, or apnea**

**Adult**

Initiate IV NSS, macro drip, KVO <sup>1</sup>  
Administer **Sodium Thiosulfate (if available):**  
(12.5 grams [50 mL] IV over 1-2 minutes <sup>2</sup>)  
Monitor ECG, if available

**Pediatric**

Initiate IV NSS, macro drip, KVO <sup>1</sup>  
Administer **Sodium Thiosulfate (if available):**  
(1.6 mL/kg, Maximum dose 12.5 grams <sup>2</sup>)  
Monitor ECG, if available

**Contact Medical Command**

**CYANIDE COMPOUND EXPOSURE  
STATEWIDE ALS PROTOCOL****Criteria:**

- A. Patients experiencing symptoms after suspected exposure to cyanide or cyanogen chloride:
  - 1. Serious exposure: symptoms include unconsciousness, seizures, and apnea. The skin may be bright red.
  - 2. Moderate exposure: symptoms may include dizziness, nausea, weakness, eye/throat irritation, and giddiness.
- B. Fire victims may be exposed to cyanide when entrapped in an enclosed structure fire. Fire victims with altered mental status, seizures, and apnea may be treated with this protocol **in addition to the Poisoning protocol #8031.**
- C. Patients exposed hydrogen sulfide in an enclosed space (for example a manure pit) that have altered mental status, seizures, or apnea may be treated with sodium thiosulfate, but a medical command physician or poison control center should be contacted before using this protocol in this situation.

**Exclusion Criteria:**

- A. Patients with suspected exposure, but without symptoms, should be evaluated for decontamination but do not require further medical treatment.
- B. If patients are seizing and have pinpoint pupils, excessive nasal/oral secretions, or muscle fasciculation (rippling tremors under skin), EMS personnel should consider exposure to nerve agents (See Nerve Agent Protocol).

**System Requirements:**

- A. Sodium thiosulfate may be carried by ALS ambulance services if the medication is permitted by the regional treatment protocols. The service must report the amount carried to the regional EMS council, and the regional EMS council should coordinate the stocks of antidote with the regional counterterrorism task forces.
- B. Until the patient has been properly decontaminated, all EMS personnel who treat patients of suspected exposure to cyanide compounds should use Level B PPE. Level B PPE should only be used by personnel with appropriate training.

**Possible Medical Command Orders:**

- A. Additional sodium thiosulfate
- B. Sodium bicarbonate for acidosis

---

**NOTES:**

- 1. In mass casualty incidents, oxygen and intravenous access should be prioritized to patients with symptoms of serious exposure if resources are limited.
- 2. May repeat sodium thiosulfate with half of initial dose once if symptoms persist after 5-10 minutes.


---

**Performance Parameters:**

- A. Every case of suspected cyanide compound exposure with any symptoms should receive QI review for appropriate use of oxygen and sodium thiosulfate.

**NERVE AGENT/PESTICIDE EXPOSURE  
STATEWIDE ALS PROTOCOL**

*Decontaminate patients* - **Contact Medical Command to order release of CHEMPACK if indicated**<sup>1</sup>  
**CAUTION: Personnel must be in appropriate PPE before treating patients who have not been decontaminated.** If possible, treat patients with severe exposure during decontamination.





### Nerve Agent Antidote Table

	<b>Adult &amp; Older Children</b> <b>&gt; 90 lbs (&gt;41 kg)</b> <b>≥ 10 y/o</b>	<b>Pediatric</b> <b>40-90 lbs</b> <b>(18-41 kg)</b> <b>4-10 y/o</b>	<b>Pediatric</b> <b>15-40 lbs</b> <b>(7-18 kg)</b> <b>6 m/o-4 y/o</b>	<b>Pediatric (Infant)</b> <b>&lt; 15 lbs</b> <b>(&lt; 7kg)</b> <b>&lt; 6 m/o</b>
<p><b><u>Moderate symptoms include:</u></b></p> <ul style="list-style-type: none"> <li>• Blurred vision</li> <li>• Excessive tearing or runny nose</li> <li>• Drooling</li> <li>• Mild shortness of breath/ wheezing</li> <li>• Vomiting</li> <li>• Diarrhea, Stomach Cramps</li> <li>• Muscle twitching or sweating at site of exposure</li> </ul>	<p>1 <b>NAAA</b> IM [atropine 2mg + pralidoxime 600 mg IM]</p>	<p>1 <b>Atropen</b> (Red) [atropine 1 mg IM]</p>	<p>1 <b>Atropen</b> (Blue) [atropine 0.5 mg IM]</p>	<p>1 <b>Atropen</b> (Yellow) [atropine 0.25 mg IM]</p>
<p><b><u>Severe symptoms include:</u></b></p> <ul style="list-style-type: none"> <li>• Altered Mental Status</li> <li>• Severe shortness of breath/ wheezing</li> <li>• General Weakness/ Severe muscle twitching</li> <li>• Incontinence (urine or feces)</li> <li>• Seizures</li> <li>• Unconsciousness</li> </ul>	<p>3 <b>NAAA(s)</b> IM [atropine 6 mg + pralidoxime 1800 mg IM]  AND Anticonvulsant 1CANA autoinjector [diazepam 10 mg IM]  OR (see box below)</p>	<p>2 <b>NAAA(s)</b> IM  OR 3 <b>Atropen</b> (Red) [atropine 3 mg IM]  AND Anticonvulsant (see box below)</p>	<p>1 <b>NAAA(s)</b> IM (if &gt; 2 y/o)  OR 3 <b>Atropen</b> (Blue) [atropine 1.5 mg IM]  AND Anticonvulsant (see box below)</p>	<p>3 <b>Atropen</b> (Yellow) [atropine 0.75 mg IM]  AND Anticonvulsant (see box below)</p>

**Adult Anticonvulsant Options:**  
(Choose one)  
Titrate until seizure stops

**Lorazepam** 1-2 mg IV/IO mg/kg, max 2 mg/dose);  
may repeat every 5 minutes until maximum of 4 mg

OR

**Diazepam** 5-10 mg IV/IO(0.01 mg/kg);  
may repeat every 5 minutes until maximum 0.3 mg/kg

OR

**Midazolam** 1-5 mg IV/IO (0.05 mg/kg);  
may repeat every 5 minutes until maximum of 0.1 mg/kg

**Pediatric Anticonvulsant Options:**  
(Choose one)  
Titrate until seizure stops

**Lorazepam** 0.1 mg/kg IV/IO/ IM (max 2 mg/dose)  
may repeat every 5 minutes until maximum of 4 mg

OR

**Diazepam** 0.3 mg/kg IV/IO/ IM  
Max 10 mg/dose IV/IO (0.5 mg/kg PR<sup>12</sup>)  
may repeat every 5 minutes until maximum of 0.6 mg/kg

OR

**Midazolam** 0.1 mg/ kg IV/IO  
Max 5 mg/dose IV/IO (0.15 mg/kg IM<sup>12</sup>)  
may repeat every 5 minutes until maximum of 0.2 mg/kg IV

**NERVE AGENT/PESTICIDE EXPOSURE  
STATEWIDE ALS PROTOCOL****CRITERIA:**

- A. Patients experiencing symptoms after suspected exposure to:  
Nerve Agents (Tabun, Sarin, Soman, VX)  
*OR*  
Organophosphate (Malathion, Parathion) / carbamate (Sevin) pesticides.
1. **Mild symptoms include:**
    - a. Pinpoint pupils
    - b. Runny nose
    - c. Suspected exposure to nerve agent, but no symptoms
  2. **Moderate symptoms include:**
    - a. Blurred vision
    - b. Excessive tearing or runny nose
    - c. Drooling
    - d. Mild shortness of breath/ wheezing
    - e. Vomiting
    - f. Diarrhea, Stomach Cramps
    - g. Muscle twitching or sweating at site of exposure
  3. **Severe symptoms include:**
    - a. Altered Mental Status
    - b. Severe shortness of breath/ wheezing
    - c. General Weakness/ Severe muscle twitching
    - d. Incontinence (urine or feces)
    - e. Seizures
    - f. Unconsciousness

**EXCLUSION CRITERIA:**

- A. Patients with suspected exposure, but without symptoms, should be decontaminated as appropriate, but do not require further medical treatment.
- B. If patients are seizing and **do not** have pinpoint pupils, excessive nasal/oral secretions, or muscle fasciculation (rippling tremors under skin), EMS personnel should consider exposure to cyanide (See Cyanide Protocol).

**SYSTEM REQUIREMENTS:**

- A. Nerve agent antidote auto-injectors (NAAAs) and pralidoxime chloride (2-PAMCl) may be carried by ALS ambulance services if the medication is permitted by the regional drug list. The service must report the amount carried to the regional EMS council, and the regional EMS council should coordinate the stocks of antidote with the regional counterterrorism task forces.
- B. Until the patient has been properly decontaminated, all EMS personnel who treat patients of suspected exposure to nerve agents should use Level B PPE. Level B PPE should only be used by personnel with appropriate training.
- C. EMTs, who have completed Department approved BLS NAAA training, may administer NAAAs under the supervision of an on-scene paramedic after the paramedic has assessed the patient and determined the number of NAAAs to be administered.
- D. BLS ambulance services may carry NAAAs for personal protection of BLS personnel. In this situation, these medications must be prescribed by the service medical director who is responsible for assuring appropriate instruction on when and how to use the medication. These NAAAs are not for patient use unless supervised by appropriate ALS personnel.

**NOTES:**

1. The Strategic National Stockpile CHEMPACKs are located at predetermined locations throughout the Commonwealth. The CHEMPACKs include autoinjectors and antidotes for nerve agent exposure. In the event of a mass casualty incident involving a suspected nerve agent, CHEMPACK(s) shall be released to an incident scene when a medical command physician orders the release of these antidotes through a county Emergency Management Agency

2. Due to severe bronchoconstriction and secretions, ventilation may be difficult, therefore atropine should be administered before attempts to intubate patient.
3. In mass casualty incidents, oxygen, intravenous access, pulse oximetry monitoring, and ECG monitoring should be prioritized to patients with severe symptoms if resources are limited.
4. NAAA (Nerve Agent Antidote Autoinjectors) are available in several brands. MARK 1 kits include 2 mg atropine and 600 mg pralidoxime in separate autoinjectors in a single kit. DuoDote autoinjectors contain 2.1 mg atropine and 600 mg pralidoxime in a single autoinjector. Atropens contain atropine in various doses depending upon the color-coded autoinjector.
5. Do not administer pralidoxime (2-PAMCI) to patients with exposure to carbamate pesticide (Sevin).
6. If NAAAs are not available, alternatively administer:
  - a. Atropine IM or IV/ IO and pralidoxime IM only, if available. Always administer atropine dose before pralidoxime dose. See Nerve Agent Antidote Table for doses.
  - b. **Mark I kits and DuoDotes are not recommended for children under 2 years old**, but appropriate Atropen or atropine doses may be given (see Nerve Agent Antidote Table).
7. **Use of the NAAAs:**
  - a. The NAAA contains either a single autoinjector or a kit with two auto injectors. These are administered IM by pressing the end of the device onto the thigh or buttocks.
    - 1) Remove the NAAA from its storage location.
    - 2) With your non-dominant hand, hold the auto injectors by the plastic clip so that the larger auto injector is on top and both are positioned in front of you at eye level.
    - 3) With the other hand check the injection site (lateral thigh muscle) for buttons or objects in the pockets which may interfere with the injections.
    - 4) Grasp the auto injector with the thumb and first two fingers. **Do not place your thumb/finger/palm over the end of the autoinjector. Atropine doses should all be administered prior to the administration of 2-PAM if using MARK 1 kits.**
    - 5) Pull the injector out of the clip with a smooth motion.
    - 6) Hold the auto injector like a pen or pencil, between the thumb and first two fingers.
    - 7) Position the green tip of the auto injector against the injection site.
    - 8) Apply firm, even pressure (not a jabbing motion) to the injector until it pushes the needle into the lateral thigh muscle.
    - 9) Hold the injector firmly in place for at least 10 seconds.
    - 10) Carefully remove the auto injector.
    - 11) Place the used auto injector into a sharps container.
    - 12) Administer additional autoinjectors using the procedures outlined in steps 4 through 11.
    - 13) Annotate the number of auto injectors administered on your patient care report or (in a mass casualty incident) on the triage tag.


---

**Performance Parameters:**

- A. Every case of suspected nerve agent or pesticide exposure with any symptoms should receive QI review for appropriate use of antidotes.

**MEDICAL COMMAND CONTACT  
STATEWIDE ALS PROTOCOL**

Follow Appropriate Protocol <sup>1,2</sup>



**MEDICAL COMMAND CONTACT  
STATEWIDE ALS PROTOCOL****Purpose of Medical Command contact:**

- A. By the Pennsylvania EMS Act and its regulations, EMS personnel will provide care within their scope of practice and will follow Department of Health-approved protocols or Medical Command orders when delivering EMS care.
- B. Medical Command must order any ALS treatment (medication or procedure) that an EMS practitioner provides when that treatment is not included in or is a deviation from the Department-approved protocols. This applies to all ALS care, including interfacility transport.
- C. In certain circumstances, as defined by the Statewide BLS Protocols, medical command must be contacted by EMS (BLS or ALS) personnel.
- D. Protocols cannot adequately address every possible patient scenario. The Pennsylvania EMS System provides a structured Medical Command system so that EMS personnel can contact a Medical Command Physician when the personnel are confronted with a situation that is not addressed by the protocols or when the EMS personnel have any doubt about the appropriate care for a patient.
- E. In some situations and geographic locations, it is not possible for an EMS practitioner to contact a medical command physician. In some protocols, there are accommodations for additional care when a medical command facility cannot be contacted.
- F. The protocol section entitled “Possible Medical Command Orders” are intended to educate EMS practitioners to the possible orders that they may receive, and as a resource to medical command physicians. Medical command physicians are not obligated to provide orders consistent with these “possible orders”. **Interventions listed under “Possible Medical Command Orders” may ONLY be done when they are ordered by a medical command physician. These possible treatments should not be done in situations where medical command cannot be contacted.**
- G. Contact with medical command may be particularly helpful in the following situations:
  - 1. Patients who are refusing treatment
  - 2. Patients with time-dependent illnesses or injuries who may benefit from transport to a specific facility with special capabilities (e.g. acute stroke, acute ST-elevation MI)
  - 3. Patients with conditions that have not responded to the usual protocol treatments.
  - 4. Patients with unusual presentations that are not addressed in protocols.
  - 5. Patients with rare illnesses or injuries that are not frequently encountered by EMS personnel.
  - 6. Patients who may benefit from uncommon treatments (e.g. unusual overdoses with specific antidotes).
- H. EMS Service Medical Directors may require more frequent contact with medical command than required by protocol for ALS personnel who have restrictions on their medical command authorization. EMS Service Medical Directors that want medical command to be contacted on every call must do this in conjunction with local medical command facilities or within a regional plan.

**Purpose of facility “EMS Notification”:**

- A. If a patient’s condition has improved and the patient is stable, interventions from a medical command physician are rarely needed, and contact with the medical command physician is disruptive to the physician’s care of other patients.
- B. When medical command is not required or necessary, regional policy may require that the receiving facility should still be notified if the patient is being transported to the Emergency Department. This “EMS notification” should be provided to the facility by phone or radio, and may be delivered to any appropriate individual at the facility.
- C. An “EMS Notification” should be a short message that includes the EMS service name or designation, the patient age/gender, the chief complaint or patient problem, and whether the patient is stable or unstable.
- D. “EMS Notification” is not necessary when a patient is not being transported to the receiving facilities Emergency Department (e.g. Inter-facility transfer to an acute care facility when the patient is a direct admission to an inpatient floor).

- E. Providing “EMS Notification” to the ED may allow a facility to be better prepared for a patient arriving by ambulance and may decrease the amount of time needed to assign an ED bed to an arriving patient.

---

**Notes:**

1. You may contact medical command regardless of your position in the protocol if you need advice or direction in caring for the patient. Medical command should be contacted for orders if a patient requiring interfacility transport needs a medication/ treatment that is not included above the contact medical command line in any Department-approved protocol.
2. When in doubt, contact medical command.
3. For example, a patient with chest pain may have almost complete resolution of pain after oxygen, aspirin, and several nitroglycerins AND may have normal vital signs.
4. Regional policy may determine the preferred method of medical command contact/ EMS notification.
5. Cellular technology may be utilized but all EMS services must maintain the ability to contact medical command by radio also.
6. **If the receiving facility is also a medical command facility, the initial medical command contact should be made to the receiving facility.** If the receiving facility cannot be contacted, an alternate facility may be contacted. The medical command physician at the alternate facility is responsible for relaying the information to the receiving facility.
7. Procedures or treatments listed after the medical command box may be considered and performed at the discretion of the ALS practitioner if unable to contact medical command if the ALS practitioner believes that these treatments are appropriate and necessary.
8. Attempts to contact medical command must be documented on the PCR, and the practitioner should document the reasons for continuing with care below the medical command box. Only mark the Medical Command section of the PA PCR if you sought Medical Command.
9. Every time medical command was contacted, the EMS practitioner must document the medical command facility, the medical command physician, and the orders received.
10. If patient condition worsens after EMS notification, contact medical command.

---

**Performance Parameters:**

- A. 100% audit of cases where treatments beyond the “contact medical command” box were performed after unsuccessful contact with medical command.
- B. Documentation of medical command facility contacted, medical command physician contacted, and orders received in every case where medical command is contacted.
- C. Review of cases for appropriate contact with medical command when required by certain protocols (e.g. acute stroke symptoms, refusal of treatment, etc...), when patient’s condition does not improve with protocol treatment, and when patients are unstable.
- D. Review of cases for appropriate use of EMS notification, and inappropriate use of medical command contact for stable patients whose symptoms were alleviated by protocol treatments.

**THIS PAGE INTENTIONALLY LEFT BLANK**

Acute Coronary Syndrome Suspected.....	5001-1 thru 5001-2
Agitated Behavior / Psychiatric Disorders.....	8001-1 thru 8001-2
Airway Management.....	4001-1 thru 4001-2
Airway Obstruction.....	3001-1 thru 3001-2
Allergic Reaction / Anaphylaxis.....	4011-1 thru 4011-2
Altered Level of Consciousness - Adult.....	7002A-1 thru 7002A-4
Altered Level of Consciousness - Pediatric.....	7002P-1 thru 7002P-4
Anaphylaxis / Allergic Reaction.....	4011-1 thru 4011-2
Asthma / COPD / Bronchospasm.....	4022-1 thru 4022-2
Asystole / Pulseless Electrical Activity (PEA) – Adult.....	3042A-1 thru 3042A-2
Asystole / Pulseless Electrical Activity (PEA) – Pediatric.....	3042P-1 thru 3042P-2
<b>Bradycardia – Adult.....</b>	<b>5021A-1 thru 5021A-2</b>
Bradycardia – Pediatric.....	5021P-1 thru 5021P-2
Bronchospasm / Asthma / COPD.....	4022-1 thru 4022-2
Burns.....	6071-1 thru 6071-4
Cardiac Arrest – Traumatic.....	3032-1 thru 3032-2
Cardiac Arrest – Hypothermia.....	3035-1 thru 3035-2
<b>Chest Pain (Suspected Acute Coronary Syndrome).....</b>	<b>5001-1 thru 5001-2</b>
Cold Injury / Hypothermia / Frostbite.....	6081-1 thru 6081-2
Confirmation of Airway Placement.....	2032-1 thru 2032-2
Congestive Heart Failure (CHF).....	5002-1 thru 5002-2
COPD / Asthma / Bronchospasm.....	4022-1 thru 4022-2
Crush Syndrome.....	6004-1 thru 6004-2
Cyanide Compound Exposure.....	8081-1 thru 8081-2
<b>Extremity Trauma.....</b>	<b>6003-1 thru 6003-2</b>
Frostbite / Hypothermia / Cold Injury.....	6081-1 thru 6081-2
<b>General Protocol Principles.....</b>	<b>1000-1 thru 1000-6</b>
Heat Emergencies.....	6086-1 thru 6086-2
Hypothermia / Cold Injury / Frostbite.....	6081-1 thru 6081-2
Major Multisystem Trauma or Traumatic Shock.....	6002-1 thru 6002-4
Medical Command Contact.....	9001-1 thru 9001-4
Narrow Complex Tachycardia – Adult.....	5022A-1 thru 5022A-2
Narrow Complex Tachycardia – Pediatric.....	5022P-1 thru 5022P-2
Nausea / Vomiting.....	7010-1 thru 7010-2
Nerve Agent / Pesticide Exposure.....	8083-1 thru 8083-4
Neonatal / Newborn Resuscitation.....	3033-1 thru 3033-2
Newborn / Neonatal Resuscitation.....	3033-1 thru 3033-2
Pesticide / Nerve Agent Exposure.....	8083-1 thru 8083-4
Psychiatric Disorders/Agitated Behavior.....	8001-1 thru 8001-2
Poisoning / Toxin Exposure.....	8031-1 thru 8031-6
Post-Partum Hemorrhage.....	7087-1 thru 7087-2
Postresuscitation Care.....	3080-1 thru 3080-2
Sedation-Assisted Intubation.....	<b>(OPTIONAL)</b> 4002-1 thru 4002-4
Seizure.....	7007-1 thru 7007-2
Seriously Ill Appearing Patient.....	7009-1 thru 7009-2
Shock.....	7005-1 thru 7005-2
<b>Suspected Acute Coronary Syndrome.....</b>	<b>5001-1 thru 5001-2</b>
Termination of Resuscitation.....	<b>(GUIDELINE)</b> 3091-1 thru 3091-2
Toxin Exposure / Poisoning.....	8031-1 thru 8031-6
Trauma (Extremity).....	6003-1 thru 6003-2
Trauma (Major Multisystem or Traumatic Shock).....	6002-1 thru 6002-4
Traumatic Shock or Major Multisystem Trauma.....	6002-1 thru 6002-4
Ventricular Fibrillation / Pulseless V Tach – Adult.....	3041A-1 thru 3041A-2
Ventricular Fibrillation / Pulseless V Tach – Pediatric.....	3041P-1 thru 3041P-2
Vomiting / Nausea.....	7010-1 thru 7010-2



Wide Complex Tachycardia – Adult.....	5023A-1 thru 5023A-2
Wide Complex Tachycardia – Pediatric.....	5023P-1 thru 5023P-2